Exam Cram: NCLEX-RN Practice Questions

Copyright © 2006 by Que Certification

International Standard Book Number: 0789732688

Warning and Disclaimer

Every effort has been made to make this book as complete and as accurate as possible, but no warranty or fitness is implied. The information provided is on an "as is" basis. The author and the publisher shall have neither liability nor responsibility to any person or entity with respect to any loss or damages arising from the information contained in this book or from the use of the CD or programs accompanying it.

When reviewing corrections, always check the print number of your book. Corrections are made to printed books with each subsequent printing. To determine the print number of your book, view the copyright page. The print number is the right-most number on the line below the "First Printing" line. For example, the following indicates that this is the 2nd printing of this title.

First Printing: July 2005

08 07 06 05 4 3 2

Corrections

Pg	Error	Correction
xvi	The allotted time is 5 hours,	The allotted time is 6 hours,
3	11client with pulmonary embolis .	client with pulmonary emboli .
3	11. A. "I will administer the medication before meals"	"I will not need to administer the medication before meals"

3	11. C. "I will check the PTT before administering the medication."	"I will not need to check the PTT before administering the medication."
3	11. D. "I will need to aspirate when I give dalteparin."	"I will avoid aspirating when I give dalteparin."
10	47following to be to a	following to be a
19	90. glyburide (diaBeta) 1.25mg	glyburide (Diabeta) 1.25mg
20	94. The client is diagnosed with multiple myoloma .	The client is diagnosed with multiple myeloma .
25	120. C. Pickles, salad with vinaigrette dressing, beef	Pickles, salad with vinigrette dressing, beef
33	160. C. A client with laryngeal cancer with a laryngetomy.	A client with laryngeal cancer with a laryngectomy.
35	169. D. "Sally has not	"She has not
37	180. C. "You will be given some medication to anesthetize the area."	"You will be asleep during the procedure."
44	215. "I'm feeing really hot."	"I'm feeling really hot."
49	238. C. Hamburger with ketchup	Hamburger with whole wheat bun
50	243. In evaluating the effectiveness of IV Pitocin for a client with secondary dystocia, the nurse should expect:	In evaluating the effectiveness of IV Oxytocin for a laboring client , the nurse should expect:
50	243. A. A rapid , painless delivery	A rapid delivery
51	248. B. Start and IV	Increase the IV infusion
51	248. C. Reposition the client	Turn the patient to the left side
51	249. A. A fetal heart rate of 170- 180bpm	A fetal heart rate of 180bpm

51	249. B. A baseline variability of 25- 35bpm	A baseline variability of 35bpm
51	249. C. Ominous periodic changes	Ominous changes
54	226. B	226. C
74	163. Answer D is incorrect because	Answer C is incorrect because
82	226. Answer B is correct.	Answer C is correct.
82	226. Answers A, C , and D are incorrect	Answers A, B , and D are incorrect
92	27Laxoxin.	Laxoxin (digoxin).
99	65prescribed imipramine for a client	prescribed Tofranil (imipramine) for a client
100	66prescribed Timoptic eyedrops.	prescribed Timoptic (timolol) eyedrops.
103	82. C. Vent contamination of the suture line	Prevent contamination of the suture line
115	147for Creon.	for Creon, a pancreatic enzyme replacement.
128	206. A client scheduled for an arthrectomy	A client scheduled for an atherectomy
143	30used to treat ulcers ,	used to treat gerd ,
149	74. Diabetes insipidus is characterized by excessive production of diluted urine.	Diabetes insipidus is characterized by excessive production of dilute urine.
155	121. Answer C is incorrect because it is an adverse reaction to Streptomycin.	Answer B is incorrect because it is an adverse reaction to Streptomycin.
156	129is usually not capable of crawling or standing.	is not capable of crawling or standing.
157	142before removing them .	before removing the tube .
165	204. Answer B is Correct. Formula for Calculation33 drops/minute Answers A, C and D are incorrect because they are inaccurate.	Answer B is Correct. Answers A, C, and D are incorrect because they are inaccurate amounts .

168	225. Answer D is correct. The client	Answer D is correct. The medication produces a burning sensation. The client
177	30. D. Spaghetti	Spaghetti with meat sauce
185	69. A client being treated with sodium warfarin has a	A client being treated with Coumadin (sodium warfarin) has a
188	80. The nurse is teaching the mother regarding treatment for enterobiasis .	The nurse is teaching the mother regarding treatment for pediculosis cupitis .
188	80. B. The entire family should be treated.	Bed linens should be washed in hot water.
188	83. The nurse caring for a client in the neonatal intensive care unit administers adult strength Digitalis to the 3-pound infant. As a result of her actions, the baby suffers permanent heart and brain damage.	The nurse caring for a client in the intensive care unit administers the wrong medication . As a result of her actions, the client suffers permanent brain damage .
189	85. The client returns to the unit from surgery with a blood pressure of 90/50, pulse 132, and respirations 30.	The client returns to the unit from surgery with hypotension tachycardia and increased respiration.
189	85. A. Continuing to monitor the vital signs	Monitor the vital signs
189	85. B. Contacting the physician	Call the physician
189	85. C. Asking the client how he feels	Explore how the client how he feels
189	85. D. Asking the LPN to continue the post op care	Asking the graduate nurse to take the vital signs
189	86. Which nurse should be assigned to care for the postpartal client with preeclampsia ?	Which nurse should be assigned to care for the postpartal client with hypertension ?
189	86. A. The RN with 2 weeks of experience in postpartum	The LPN with 2 weeks of experience in postpartum

189	86. B. The RN with 3 years of experience in labor and delivery	The RN with 1 years of experience in labor and delivery
189	86. C. The RN with 10 years of experience in surgery	The RN with 10 years of experience in medical nursing
189	86. D. The RN with 1 years of experience in neonatal intensive care unit	The RN with 3 years of experience in neonatal intensive care unit
190	91. The nurse is caring for a 6-year old client admitted with a diagnosis of conjunctivitis. Before administering eyedrops, the nurse	Before administering eyedrops to a child , the nurse
190	91. A. The eye should be cleansed with warm water, removing any exudate, before instilling the eyedrops.	The eyelid should be cleansed before instilling the eyedrops.
190	91. B. The child should be allowed to instill his own eyedrops.	The child should instill his own eyedrops.
190	91. C. The mother should be allowed to instill the eyedrops.	The mother should not be allowed to instill the eyedrops.
190	91. D. If the eye is clear from any redness or edema, the eyedrops should be held.	The eyedrops should be administered inner to outer canthas.
192	101. A gravida III para 0 is admitted to the labor and delivery unit. The doctor performs an amniotomy.	Delete both sentences.
192	101. A. Fetal heart tones 160bpm	Tachycardia
192	101. B. A moderate amount of straw-colored fluid	Straw-colored fluid
192	101. C. A small amount of greenish fluid	Greenish fluid

192	101. D. A small amount of the umbilical cord	The umbilical cord
192	102. The client is admitted to the unit. A vaginal exam reveals that she is 2em dilated. Which of the following statements would the nurse expect her to make?	Which of the following statements would the nurse expect the laboring client during the active phase of labor to make?
193	103. B. Apply the fetal heart monitor	Apply an internal fetal heart monitor
193	103. D. Perform an ultrasound exam	Perform an ultrasound
193	108. As the client reaches 8cm dilation,	As the client reaches 6cm dilation,
193	108. The FHR baseline is 165-175bpm with variability of 0-2bpm.	Delete sentence.
193	108. A. The baby is asleep .	The baby is sleeping
193	108. C. There is a vagal response .	There is a head compression .
204	159. D. Bromocrystine (Pardel)	Bromocriptine (Parlodel)
210	189. A client is admitted for an MRI.	A client is admitted for an CAT scan .
212	201. A client with pneumacystis carini pneumonia is receiving trimetrexate .	A client with pneumacystis carini pneumonia is receiving methotrexate .
219	239. The infant is admitted to the unit with tetrology of falot .	The infant is admitted to the unit with tetralogy of fallot .
224	41. B	A
225	169. B	D

232	41. Answer B is correct. Proton pump inhibitors such as Nexium and Protonix should be taken with meals, for optimal effect. Histamine-blocking agents such as Zantac should be taker 30 minutes before meals, so answer A is incorrect.	as Nexium and Protonix should be taken 30 minutes prior to meals, for optimal effect.
234	65. The client might experience symptoms of hypoxia if he becomes dehydrated or deoxygenated; extreme exercise, especially in warm weather, can exacerbate the condition.	Delete sentence.
235	67at this time and could cause hyponatremia and further hypotension.	at this time.
236	80. Entire entry deleted and rewritten.	Answer B is correct. Pediculosis capitis or head lice is treated with Kwell shampoo. Bed linen should be washed in hot water to assure that all lice and eggs are killed. A is incorrect because medication can be used on younger children. C is incorrect because toxicity can occur with repeated use. D is not true.
237	83. Giving the infant an overdose falls into this category.	Giving the wrong medication falls into this category.
237	86. The nurse with 3 years of experience in	The nurse with 2 years of experience in

238	91. A 6-year old child is not developmentally ready to instill his own eyedrops, so answer B is incorrect. Although the mother of the child can instill the eyedrops, the area must be cleansed before administration, making answer C incorrect. Although the eye might appear to be clear, the nurse should instill the eyedrops as ordered, so answer D is incorrect.	A child is not developmentally ready to instill his own eyedrops, so answer B is incorrect. Leaves as is disregard line, making answer C incorrect. Oinment is instilled inner to outer canthas not drops, so D is wrong.
239	102. Dilation of 2cm marks the end of the latent phase of labor.	The client can get her epidural during the active phase of labor.
247	159magnesium sulfate, and Pardel is a	magnesium sulfate, and Parlodel is a
247	159. Pardel was used at one time to dry breas milk.	Parlodel was used at one time to dry breast milk.
248	169. Answer B is correct. Hypospadia is a condition in which there is an opening on the dorsal side of the penis. Answer A is incorrect because hypospadia does not concern the urethral opening. Answer C is incorrect because the size of the penis is not affected. Answer D is incorrect because the opening is on the dorsal side , not the vertical side.	Answer D is correct. Hypospadia is a condition in which there is an opening on the vertical side of the penis. Answer A is incorrect because hypospadia does involves an abnormal urethral opening. Answer C is incorrect because the size of the penis is not affected. Answer B is incorrect because the opening is on the vertical side, not the dorsal side.
251	189not have an MRI because	not have a CAT scan because
251	189hip replacement can have an MRI, so	hip replacement can have an MRI or CAT scan, so
257	239. The infant with tetrology of falot has	The infant with tetralogy of fallot has

272	67. A. Decreased blood pressure and papillary meiosis	Decreased blood pressure and pupillary meiosis
280	106. A. 10cc	10ml
280	106. B. 11.5cc	11.5ml
280	106. D. 11.5g	12gm
280	108. Which position is best for the newborn with RDS?	Which position is best for maintaining an open airway?
282	115urine is very diluted .	urine is very dilute .
291	162irrigation to a client	irrigation for a client
315	21is diagnostic for fetal wall defects , most notably neural tube defects.	is diagnostic for neural tube defects.
318	43. Answers A and B are incorrect because	Answers A and C are incorrect because
319	49client with gallstones.	client with stones in the gall bladder.
321	67and papillary meiosis.	and pupillary meiosis.
327	106for urine output of 11.5g .	for urine output of 11.5ml .
348	5. B. (Nalonxone) Naloxone	(Nalonxone) Narcan
388	198. B. Naloxonel (Narcan)	Naloxone (Narcan)
391	214. B. Urine output of 30 cc /hr	urine output of 30 mL /hr
391	215reveal a phenytoin level of 38mcg/mL.	reveal a theophylline level of 38mcg/mL.
408	45. Assessing for Tinel's syndrome is	Assessing for Tinel's sign is
409	55. The client is exhibiting symptoms of air embolism.	The client is at risk for air embolism.

412	75failure that happen commonly	failure that occurs commonly
412	82. Pain medications are contraindicated in neurological clients, so	Pain medications are contraindicated in undiagnosed neurological clients, so
414	10050% BSA=13,600 cc	50% BSA=13,600 mL
417	121. Chest tube removal prevents air entrance during the procedure.	During chest tube removal, this procedure prevents air entrance into the chest cavity.
422	171identify the pulse preoperatively	identify the pulses preoperatively
424	184. The highest calcium level is in the granola and grapefruit.	The highest calcium level is in the bagel and skim milk.
426	204. Vomiting, a heart rate 120,	Vomiting, a heart rate of 120,

This errata sheet is intended to provide updated technical information. Spelling and grammar misprints are updated during the reprint process, but are not listed on this errata sheet.