

# EXAM ✓ CRAM

Succeed with topical reviews, practice exams, and preparation tools

# NCLEX-PN<sup>®</sup>

SECOND EDITION

EXAM ✓ CRAM



The Global Leader  
in Education Publishing



Wilda Rinehart  
Diann Sloan  
Clara Hurd

## **NCLEX-PN® Exam Cram, Second Edition**

### **Copyright © 2008 by Pearson Education**

All rights reserved. No part of this book shall be reproduced, stored in a retrieval system, or transmitted by any means, electronic, mechanical, photocopying, recording, or otherwise, without written permission from the publisher. No patent liability is assumed with respect to the use of the information contained herein. Although every precaution has been taken in the preparation of this book, the publisher and author assume no responsibility for errors or omissions. Nor is any liability assumed for damages resulting from the use of the information contained herein.

ISBN-13: 978-0-7897-2706-9

ISBN-10: 0-7897-3706-x

Library of Congress Cataloging-in-Publication Data

Rinehart, Wilda.

NCLEX-PN exam cram / Wilda Rinehart, Diann Sloan, Clara Hurd. -- 2nd ed.

p. cm.

ISBN 978-0-7897-3706-9 (pbk. w/cd)

1. Practical nursing--Examinations, questions, etc. 2. Nursing--Examinations, questions, etc. 3. National Council Licensure Examination for Practical/Vocational Nurses--Study guides. I. Sloan, Diann. II. Hurd, Clara. III. Title.

RT62.R55 2008

610.73'076--dc22

2008000133

Printed in the United States of America

First Printing: February 2008

### **Trademarks**

All terms mentioned in this book that are known to be trademarks or service marks have been appropriately capitalized. Pearson Education cannot attest to the accuracy of this information. Use of a term in this book should not be regarded as affecting the validity of any trademark or service mark.

### **Warning and Disclaimer**

Every effort has been made to make this book as complete and as accurate as possible, but no warranty or fitness is implied. The information provided is on an "as is" basis. The authors and the publisher shall have neither liability nor responsibility to any person or entity with respect to any loss or damages arising from the information contained in this book or from the use of the CD or programs accompanying it.

### **Bulk Sales**

Que Publishing offers excellent discounts on this book when ordered in quantity for bulk purchases or special sales. For more information, please contact

**U.S. Corporate and Government Sales**

**1-800-382-3419**

**corpsales@pearsontechgroup.com**

For sales outside of the U.S., please contact

**International Sales**

**international@pearsoned.com**

### **Publisher**

Paul Boger

### **Associate Publisher**

David Dusthimer

### **Acquisitions Editor**

Betsy Brown

### **Senior Development Editor**

Christopher Cleveland

### **Managing Editor**

Patrick Kanouse

### **Senior Project Editor**

San Dee Phillips

### **Indexer**

Ken Johnson

### **Technical Editor**

Kathy Heston

Mercy Heston

### **Publishing Coordinator**

Vanessa Evans

### **Book Designer**

Gary Adair

### **Page Layout**

Mark Shirar

# Introduction

## Welcome to the NCLEX-PN<sup>®</sup> Exam Cram

This book will help you prepare to take and pass the Licensure Exam for Practical Nurses. This Introduction discusses the NCLEX<sup>®</sup> exam in general and how the *Exam Cram* can help you prepare for the test. It doesn't matter whether this is the first time you're going to take the exam or if you have taken it previously; this book gives you the necessary information and techniques to obtain licensure.

*Exam Cram* books help you understand and appreciate the subjects and materials you need to pass. The books are aimed at test preparation and review. They do not teach you everything you need to know about the subject of nursing. Instead they present materials you are likely to encounter on the exam.

Using a simple approach, we help you understand the need-to-know information. First, you learn content as it applies to medical-surgical nursing, psychiatric-mental health nursing, obstetric nursing, and pediatric nursing, with an emphasis on pharmacology, skills, and management of these disorders. In a well-organized format, you learn the pathophysiology of the most common problems affecting clients, the treatment of these disorders, and the nursing care required.

The NCLEX-PN<sup>®</sup> consists of questions from the cognitive levels of knowledge, comprehension, application, and analysis. The majority of questions are written at the application and analysis levels. Questions incorporate the five stages of the nursing process (assessment, diagnosis, planning, implementation, and evaluation) and the four categories of client needs. Client needs are divided into subcategories that define the content within each of the four major categories. These categories and subcategories are

- ▶ A. Safe, effective care environment:
  - ▶ Coordinated care: 12%–18%
  - ▶ Safety and infection control: 8%–14%
- ▶ B. Health promotion and maintenance: 7%–13%
- ▶ C. Psychosocial integrity: 8%–14%

- ▶ D. Physiological integrity:
  - ▶ Basic care and comfort: 11%–17%
  - ▶ Pharmacological and parenteral therapy: 9%–15%
  - ▶ Reduction of risk: 10%–16%
  - ▶ Physiological adaptation: 11%–17%

## Taking the Computerized Adaptive Test

Computer Adaptive Testing offers the candidate several advantages. The graduate can schedule the exam at a time that is convenient for him. The Pearson VUE testing group is responsible for administering the exam. Because you might not be familiar with the Pearson VUE testing centers, we recommend that you arrive at least 30 minutes early to acclimate yourself to the surroundings and learn what you need to do while testing at the center. If you are late, you will not be allowed to test. Bring two forms of identification with you, one of which must be a picture ID. Be sure that your form of identification matches your application. You will be photographed and fingerprinted upon entering the testing site, so don't let this increase your stress. The allotted time is 5 hours. The candidate can receive results within approximately 7 days (in some states even sooner). Remember that the exam is written at approximately the 10th-grade reading level so keep a good dictionary handy during your studies.

## The Cost of the Exam

The candidate wanting to take the licensure exam must fill out two applications, one to the National Council and one to the state in which she wants to be licensed. A separate fee must accompany each application. The fee required by the National Council is \$200. State licensing fees vary from state to state. The candidate should contact the state where she wishes to become licensed for a list of fees for that specific state. Licensure applications can be obtained on the National Council's website at [www.ncsbn.org](http://www.ncsbn.org). Several states are members of the multistate licensure compact. This means that, if you are issued a multistate license, you pay only one fee. This information can also be obtained by visiting the National Council's website. A list of phone numbers and websites is included on this book's CD, in Appendix C, "Alphabetical Listing of Nursing Boards in the United States and Protectorates."

# How to Prepare for the Exam

Judicious use of this book, either alone or with a review seminar, such as that provided by Rinehart and Associates, will help you to achieve your goal of becoming a practical nurse. As you review for the NCLEX® Exam, we suggest that you find a location where you can concentrate on the material each day. A minimum of 2 hours per day for at least 2 weeks is suggested. We have provided you with exam alerts, tips, notes, and sample questions, both multiple-choice and alternative items. These questions will acquaint you with the type of questions you will see during the exam. We have also formulated a mock exam, with those difficult management and delegation questions, which you can score to determine your readiness to test. Pay particular attention to the Exam Alerts and the Cram Sheet. Using these will help you gain and retain knowledge and help reduce your stress as you prepare to test.

# How to Use This Book

Each topical *Exam Cram* chapter follows a regular structure and includes cues about important or useful information. Here's the structure of a typical chapter:

- ▶ **Opening hotlists**—Each chapter begins with a list of terms and concepts you must learn and understand before you can know the subject matter. The hotlists are followed by an introductory section to set the stage for the rest of the chapter.
- ▶ **Topical coverage**—After the opening hotlists, each chapter covers a series of topics related to the chapter's subject title.

Even though the book is structured to the exam, these flagged items are often particularly important:

- ▶ **Exam Alert**—Exam alerts normally stress concepts, terms, or activities that are related to one or more test questions. Anything found in exam alert format is worthy of greater attention on your part. This is what an exam alert looks like:

## CAUTION

Exam alerts are provided as a heads up that the content mentioned here might appear on the NCLEX-PN® exam.

- ▶ **Notes**—Throughout each chapter additional information is provided that, although not directly related to the exam itself, is still useful and will aid your preparation. A sample note is shown here:

**NOTE**

This is how notes are formatted. Notes direct your attention to important pieces of information that relate to nursing and nursing certification.

- ▶ **Tips**—A tip might tell you another way of accomplishing something in a more efficient or time-saving manner. An example of a tip is shown here:

**TIP**

This is how tips are formatted. Keep your eyes open for these, and you'll learn some interesting nursing tips!

- ▶ **Exam Prep Questions**—Although we talk about test questions and topics throughout the book, the section at the end of each chapter presents a series of mock test questions and explanations of both correct and incorrect answers.
- ▶ **Practice Exams**—This book offers two exams written in the NCLEX<sup>®</sup> format. These have been provided to help you evaluate your readiness to test. Answers and rationale to these questions have also been provided. We suggest that you score the exam by subtracting the missed items from the total and dividing the total answered correctly by the total number of questions. This will give you the percentage of answers correctly. We suggest that you achieve a score of at least 77% before you schedule your exam.
- ▶ **Glossary**—At the end of the book is a glossary that defines critical nursing terms we cover in this book.
- ▶ **The CD**—The CD includes a testing engine with many practice questions that you should use repeatedly to practice your test-taking skills and measure your level of learning. You should be able to correctly answer more than 77% of the questions on the practice tests before trying the real exam. The CD also contains Appendix A, “Things You Forgot”; Appendix B, “Need to Know More?”; and Appendix C, “Alphabetical Listing of Nursing Boards in the United States and Protectorates.”

- ▶ **Cram Sheet**—At the beginning of the book is a tear card we call the Cram Sheet. This is a helpful tool that gives you distilled, compressed facts and is a great tool for last-minute study and review.

## About the Book

The topics in this book have been structured using the systems approach to nursing. We believe that a simple approach to learning the disease process, treatments, and diagnostic studies is best. We review material related to diseases of each body system; the related nursing skills; and the diagnostic tests, nutrition, and pharmacology associated with each. We also consider cultural and religious aspects as they relate to the care of clients with specific illnesses.

Aside from being a test preparation book, this book is also useful if you are brushing up on your nursing knowledge. It is an excellent quick reference for the licensed nurse.

## Contact the Authors

The authors of this text are interested in you and want you to pass on the first attempt. If, after reviewing with this text, you would like to contact the authors, you can do so at Rinehart and Associates, PO Box 124, Booneville, MS 38829 or by visiting our website at [www.nclexreview.net](http://www.nclexreview.net). You can also contact us by phone at 662-728-4622.

# 7

## CHAPTER SEVEN

# Caring for the Client with Burns

---

### Terms you'll need to understand:

- ✓ Allograft
- ✓ Autograft
- ✓ Biosynthetic graft
- ✓ Burn shock
- ✓ Consensus formula
- ✓ Contracture
- ✓ Debridement
- ✓ Donor site
- ✓ Emergent phase of burn injury
- ✓ Eschar
- ✓ Heterograft
- ✓ Homograft
- ✓ Intermediate phase of burn injury
- ✓ Jobst garment
- ✓ Lund and Browder method
- ✓ Palm method
- ✓ Parkland formula
- ✓ Rehabilitative phase of burn injury
- ✓ Rule of Nines
- ✓ Total body surface area (TBSA)

---

### Nursing skills you'll need to master:

- ✓ Performing sterile dressing change
- ✓ Administering medications
- ✓ Transfusing blood and blood products
- ✓ Performing tracheostomy suction and care
- ✓ Monitoring central venous pressure
- ✓ Caring for central lines
- ✓ Assessing a burn injury using the Rule of Nines
- ✓ Calculation of IV fluid requirements using the Parkland formula and the Consensus formula



Although the incidence of burn injury has declined, burns still account for about 2,000,000 injuries each year in the United States. According to the American Burn Association (2000), more than 51,000 persons require hospital care each year for treatment of their injuries. Those with burns greater than 25% total body surface area (TBSA) are at risk of dying from smoke inhalation and other complications associated with burns. Young children and the elderly are particularly vulnerable to local and systemic effects of burns because their skin is naturally thinner. Burns are the third leading cause of death in children under age 14 and are in the top 10 of causes of death for all age groups.

Burns generally occur from one of three major sources:

- ▶ Thermal injuries (hot liquid, open flame)
- ▶ Electrical injuries (household current, lightning)
- ▶ Chemical injuries (alkaline or acid liquids or powders)

Radiation injuries are most likely to occur with industrial accidents where radioactive energy is produced or in situations where radioactive isotopes are used. More discussion on radiation injuries can be found in Chapter 18, “Emergency Nursing.”

Most burns are thermal injuries that occur in the home. Cooking accidents from hot grease or stove fires result in a significant number of injuries, as do scalds from bath water that is too hot.

### CAUTION

To prevent burns, hot water heaters should be set no higher than 120° Fahrenheit.

Carbon monoxide, sulfur oxides, cyanide, chlorine, and other toxins are released from household contents during a fire. Inhalation of these gases damages the lower airway, resulting in the collapse of the alveoli and increasing the possibility of acute respiratory distress syndrome.

## Burn Classifications

Before discussing caring for the client with burns, we must first look at how burns are classified. Treatment of the client with burns is dictated by whether the injury is classified as a *minor burn*, *moderate burn*, or *major burn*. These classifications are dependent on the degree of tissue involved and the total body surface area affected by the injury. Burns are further classified in terms of the depth of tissue destroyed or the *thickness* of the burn injury. The following list gives you an idea of the different degrees of burns, the symptoms experienced with the injury, and the expected time of healing:

- ▶ **Superficial partial thickness (first degree)**—Tissue damage is confined to the epidermis and possibly a portion of the dermis. This is the type of injury produced by sunburn or a low-intensity flash. The skin appears red but blanches with pressure. Blisters may or may not be present. The client usually complains of tingling, increased skin sensitivity, and pain that is relieved by the application of cool water or lotions containing aloe. The injury heals within a week. Although the skin peels, there is no scarring.
- ▶ **Deep partial thickness (second degree)**—Tissue damage involves the epidermis, upper dermis, and portions of the deeper dermis. Deep partial thickness injury is common in scalds and flash flames. The area involved appears blistered with weeping and edema. The client experiences pain and increased skin sensitivity, which increases with exposure to air. The use of sterile sheets and overbed cradles minimizes contact with the air and makes the client more comfortable. Morphine sulfate or other opiate analgesics are given intravenously to control pain.

**CAUTION**

Pain medication is given intravenously to provide quick, optimal relief and to prevent overmedication as edema subsides and fluid shift is resolving.

Deep partial thickness injury generally heals in two to four weeks, although infection can delay healing. Infection can also take a deep partial thickness injury to a full thickness injury.

- ▶ **Full thickness (third degree)**—Tissue damage involves the epidermis and entire dermis. The damage usually extends into subcutaneous tissue, including connective tissue, muscle, and bone. Full thickness burns result from prolonged exposure to hot liquids or open flame, electrical current, or exposure to chemical agents. Depending on the source of the injury, the affected area can appear dry, pale white, edematous, leathery, or charred. Destruction of nerve endings leaves the affected areas relatively pain free. Complicating the care of the client with full thickness injury is the development of hypovolemic burn shock, hyperkalemia, and anemia. Electrical injuries, which appear as whitish areas at the points of entry and exit, can result in changes in heart rhythm or complete cardiac standstill.

**CAUTION**

The cardiac status of a client with electrical burns should be closely monitored for at least 24 hours following the injury to detect changes in electrical conduction of the heart.


**CAUTION**

Full thickness burns can damage muscles, leading to the development of myoglobinuria, in which urinary output becomes burgundy in color. The client with myoglobinuria may require hemodialysis to prevent tubular necrosis and acute renal failure.

## Burn Measurement with TBSA

A second means of classifying burns is based on the percentage of tissue injured. Three methods are used to determine the total body surface area injured in a burn:

- ▶ **The Rule of Nines**—The Rule of Nines assigns percentages of 9 to major body surfaces. The breakdown is as follows: head = 9%, anterior trunk = 18%, posterior trunk = 18%, arms = 9% each, legs = 18% each, and perineum = 1%. The rule is demonstrated in Figure 7.1.



**FIGURE 7.1** The Rule of Nines.

- ▶ **Lund and Browder method**—The Lund and Browder method of determining TBSA is more precise because it takes into account that anatomic parts, especially the head and legs, change with growth. Special charts divide the body into very small parts and provide for an estimate of the proportion of TBSA burned. The Lund and Browder method is used to estimate TBSA in children.

- ▶ **The palm method**—The percentage affected by scattered burns may best be calculated using the palm method. The size of the client's palm represents approximately 1% of the TBSA.

Minor burn injury involves a second degree burn or less than 15% of TBSA in adults and less than 10% in children. Or, it can involve a third degree burn of less than 2% TBSA but not involving areas requiring special care (face, eyes, ears, perineum, and joints of hands and feet). Minor burns do not include electrical burn injury, inhalation injury, those clients with concurrent illness or trauma, or age-related considerations.

Moderate burn injury involves second degree burns of 15%–20% TBSA in adults, 10%–20% in children, or third degree burns less than 10% TBSA that do not involve special care areas. Moderate burns, like minor burns, do not include electrical or inhalation injury, nor those with concurrent illness, trauma, or age-related considerations.

Major burn injury involves second degree burns greater than 25% TBSA in adults, 20% in children, or all third degree burns greater than 10% TBSA. Major burns include all burns involving the structures of the head and face, hands, feet, and perineum as well as electrical and inhalation injury, concurrent illness, and trauma regardless of age.

### CAUTION

It will be beneficial to review your nursing textbooks for local and systemic reactions to burns because these injuries affect all body systems and cardiovascular and renal function in particular.

## Nursing Care for Burn Victims

Caring for a burned client represents a unique challenge to even the most experienced nursing staff because few injuries pose a greater threat to the client's physical and emotional well-being. There are three phases of burn injury, each requiring various levels of client care. The three phases are

- ▶ Emergent
- ▶ Intermediate
- ▶ Rehabilitative

## Psychological Care of a Burn Patient

Although interventions are focused on meeting the client's physiological needs during the emergent period, the nurse should keep in mind that the nature of the injury represents a time of extreme crisis for both the client and his family. Every effort should be made to provide emotional support by providing understandable explanations of procedures and making sure that the client is kept as comfortable as possible. When necessary, appropriate referrals should be made to clergy and other professionals. Interventions directed at stabilizing the client's condition as well as the type of emotional support will change as the client moves through the emergent, intermediate, and rehabilitative phases of injury.

## The Emergent Phase

The emergent phase begins with the onset of burn injury and lasts until the completion of fluid resuscitation or a period of about the first 24 hours. During the emergent phase, the priority of client care involves maintaining an adequate airway and treating the client for burn shock.

Emergency care of burns at the site of injury includes

- ▶ Extinguishing the burn source
- ▶ Soaking the burn with cool water to relieve pain and to limit local tissue edema
- ▶ Removing jewelry and nonadherent clothing
- ▶ Covering the wound with a sterile (or at least clean) dressing to minimize bacterial contamination
- ▶ Brushing off chemical contaminants, removing contaminated clothing, and flushing the area with running water

### CAUTION

The eyes should be irrigated with water immediately if a chemical burn occurs. Follow-up care with an ophthalmologist is important because burns of the eyes can result in corneal ulceration and blindness.

## Major Burns in the Emergent Phase

If the injury is determined to be a major burn injury, the following additional interventions will be taken during the emergent phase of burn care. Assessment of the following needs to take place during this phase:

- ▶ Airway
- ▶ Breathing
- ▶ Circulation

**CAUTION**

Important steps in treating a burn client include

- ▶ **Treat airway and breathing**—Traces of carbon around the mouth or nose, blisters in the roof of the mouth, or the presence of respiratory stridor indicate the client has respiratory damage. Endotracheal intubation with assisted ventilation might be required to achieve adequate oxygenation.
- ▶ **Ensure proper circulation**—Compromised circulation is evident by slowed capillary refill, a drop in normal blood pressure, and decreased urinary output. These symptoms signal impending burn shock.

These interventions come next:

- ▶ Insertion of a large bore catheter for administering IV fluids
- ▶ Calculation of TBSA involved
- ▶ Calculation of fluid needs according to one of the fluid resuscitation formulas

**CAUTION**

It is important to remember that the actual burns might not be the biggest survival issue facing burn clients. Carbon monoxide from inhaled smoke can develop into a critical problem as well. Carbon monoxide combines with hemoglobin to form carboxyhemoglobin, which binds to available hemoglobin 200 times more readily than with oxygen. Carbon monoxide poisoning causes a vasodilating effect, making the client have a characteristic cherry red appearance. Interventions for carbon monoxide poisoning focus on early intubation and mechanical ventilation with 100% oxygen.

In the hours immediately following a major burn injury, loss of capillary permeability allows intravascular fluid to flood into the extracellular space. During the emergent or resuscitative phase, efforts are directed at preventing or reversing burn shock using fluid replacement formulas. Although there are a number of acceptable formulas for calculating fluid requirements, the Parkland formula and Consensus formula are most often used.

**The Parkland Formula**

The Parkland formula provides a large volume of IV fluid in the first 24 hours to prevent deepening hypovolemic shock and further acidosis. After the first 24 hours, the amount of fluid infused should be titrated according to the urinary output, with the goal of maintaining the output between 30 ml and 50 ml per hour.

The following example steps you through a calculation of TBSA using the Rule of Nines and the fluid requirements using the Parkland formula:

A client receives full thickness burns of the arms, chest, back, and head at 0600 hours. The client weighs 180 pounds. Using the Parkland formula, how much fluid should the client receive by 1400?

**Parkland formula:**

*Ringer's Lactate*  $4 \text{ ml} \times \text{kg body weight} \times \% \text{ TBSA}$

Half of the amount is to be infused in the first 8 hours.

The remainder is to be infused over the next 16 hours.

With this information, what steps should you follow? The steps given below will help you calculate this if you have difficulty:

1. Calculate the TBSA using the Rule of Nines:

$$\text{arms (9\% each arm)} = 18\% + \text{chest (18\%)} + \text{back (18\%)} + \text{head (9\%)} = 63\%$$

2. Convert the client's weight from pounds to kilograms:

$$180 \text{ pounds} \div 2.2 \text{ pounds (2.2 pounds = 1 kg)} = 81.8 \text{ kg (round to 82 kg)}$$

3. Calculate using the Parkland formula for fluid resuscitation:

$$4 \text{ ml} \times 82 \text{ kg} \times 63 = 20,664 \text{ ml in 24 hours}$$

According to the Parkland formula, half the calculated volume of Lactated Ringer's solution is to infuse in the first 8 hours; one fourth is to infuse in the second 8 hours; and one fourth is to infuse in the remaining 8 hours.

4. The injury occurred at 0600; the first 8 hours will end at 1400. Therefore, the client should receive one half the total amount or 10,332 ml.

**The Consensus Formula**

Here's how you use the Consensus formula (for comparison with use of the Parkland formula):

**Consensus formula:**

*Ringer's Lactate or other balanced saline solution*  $2 \text{ ml} - 4 \text{ ml} \times \text{kg body weight} \times \% \text{ TBSA}$

Half of the amount is to be infused over the first 8 hours.

The remainder of the amount is to be infused over the next 16 hours.

**CAUTION**

Fluid replacement formulas are calculated from the time of injury rather than from the time of arrival in the emergency room.

With this information, what steps should you follow? The steps given here will help you calculate this if you have difficulty:

1. Calculate the TBSA using the Rule of Nines:

arms (9% each arm) = 18% + chest (18%) + back (18%) + head (9%) = 63%

2. Convert the client's weight from pounds to kilograms:

180 pounds  $\div$  2.2 pounds (2.2 pounds = 1 kg) = 81.8 kg (rounded to 82 kg)

3. Calculate using the Consensus formula for fluid resuscitation:

$2 \text{ ml} \times 82 \times 63 = 10,332 \text{ ml}$

$4 \text{ ml} \times 82 \times 63 = 20,664 \text{ ml}$

On the low end (2 ml), the amount to infuse over 24 hours would be 10,332 ml, with half to be infused in the first 8 hours and the remainder to be infused over the next 16 hours.

On the high end (4 ml), the amount to infuse over 24 hours would be 20,664 ml, with half to be infused in the first 8 hours and the remainder to be infused over the next 16 hours.

## Additional Interventions

These additional interventions are taken after assessment of airway and establishing IV access for fluid replacement. Airway and maintaining fluid volume take priority over all the other interventions:

- ▶ Administering a tetanus booster
- ▶ Inserting a urinary catheter for determining hourly output
- ▶ Inserting a nasogastric tube attached to low suction to minimize aspiration

### NOTE

Enteral feedings are usually instituted within the first 24 hours to meet the client's increased caloric needs and maintain the integrity of the intestinal mucosa thereby minimizing systemic sepsis.

- ▶ Elevating burned extremities to lessen edema formation



## The Intermediate Phase

The intermediate phase of burn care begins about 48–72 hours following the burn injury. Changes in capillary permeability and a return of osmotic pressure bring about diuresis or increased urinary output. If renal and cardiac functions do not return to normal, the added fluid volume, which prevented hypovolemic shock, might now produce symptoms of congestive heart failure. Assessment of central venous pressure provides information regarding the client's fluid status.

### NOTE

The central venous pressure (CVP) is read with the client in a supine position with the manometer level with the fourth intercostal space midaxillary line (often referred to as the *phlebostatic axis*). The normal CVP varies but the general range is between 5–12 mm H<sub>2</sub>O. Increased CVP indicates fluid volume overload; decreased CVP indicates fluid volume deficit.

Additional complications found during the intermediate phase include infections, the development of Curling's ulcer, paralytic ileus, anemia, disseminated intravascular coagulation, and acute respiratory failure.

### NOTE

Infections represent a major threat to the post-burn client. Bacterial infections (*staphylococcus*, *proteus*, *pseudomonas*, *escherichia coli*, and *klebsiella*) are common due to optimal growth conditions posed by the burn wound; however, the primary source of infection appears to be the client's own intestinal tract. As a rule, systemic antibiotics are avoided unless an actual infection exists.

During the intermediate phase, attention is given to removing the eschar and other cellular debris from the burned area. *Debridement*, the process of removing eschar, can be done placing the client in a tub or shower and gently washing the burned tissue away with mild soap and water or by the use of enzymes, substances that digest the burned tissue. Santyl (collagenase) is an important debriding agent for burn wounds.

### CAUTION

Enzymatic debridement should not be used for burns greater than 10% TBSA, for burns near the eyes, or for burns involving muscle.

Following debridement, the wound is treated with a topical antibiotic and a dressing is applied (more on dressings is covered in the next section). Commonly used topical antibiotics include

silver sulfadiazine (Silvadene); mafenide acetate (Sulfamylon); and silver nitrate, which can be used in an aqueous solution of 0.5% or Acticoat, a prepared dressing impregnated with silver nitrate. Silver nitrate has bacteriostatic properties that inhibit bacterial growth. Mafenide acetate, although painful, is useful in preventing *Pseudomonas* infections. Silvadene cools and soothes the burn wound but does not prevent infection.

## Dressings for Burns

Dressings for burns include standard wound dressings (sterile gauze) and biologic or biosynthetic dressings (grafts, amniotic membranes, cultured skin, and artificial skin).

### Standard Wound Dressings

The use of standard wound dressings makes the client more comfortable by preventing exposure of the wound to air. These dressings are usually applied every shift or once a day.

### Biologic or Biosynthetic Dressings

Biologic dressings are obtained from either human tissue (homograft or allograft) or animal tissue (heterograft or xenograft). These dressings, which are temporary, are used for clients with partial thickness or granulating full thickness injuries. The type of biologic dressing used depends on the type of wound and availability of the graft.

Homografts or allografts are taken from cadaver donors and obtained through a skin bank. These grafts are expensive and there is a risk of blood-borne infection. Heterografts or xenografts are taken from animal sources. The most common heterograft is pigskin because of its compatibility with human skin.

#### CAUTION

Certain religious and ethnic groups would be offended if offered a porcine (pigskin) graft.

Amniotic membrane is used for full thickness burns because it adheres immediately to the wound. It is also an effective covering for partial thickness burns until reepithelialization occurs. Amniotic membrane is low in cost, and its size allows for coverage of large wounds.

Cultured skin can be obtained by using a biopsy of epidermal cells taken from unburned portions of the client's body. The cells are grown in a laboratory and grafted to generate permanent skin. The process is long and costly, and extreme care is needed to prevent damage and loss of the graft.

Artificial skin (Integra) made of synthetic material and animal collagen becomes a part of the client's skin. The graft site is pliable, there is less hypertrophic scarring, and its use is helping

to eliminate the need for compression dressings like the Jobst garment during the rehabilitative phase of care.

Permanent grafts include the autograft or skin transferred from an unburned area of the client's body to the burn wound. The client generally experiences more pain from the donor site than from the burn wound because the donor site has many pain receptors. The client should receive pain medication, and both the donor site and graft site should be carefully monitored for signs of infection.

## The Rehabilitative Phase

The last stage in caring for a client with burn injury is the rehabilitative stage. Technically, this stage begins with closure of the burn and ends when the client has reached the optimal level of functioning. In actuality, it begins the day the client enters the hospital and can continue for a lifetime. In the emergent and intermediate phases, the focus is on establishing and maintaining physiological equilibrium. In the rehabilitative phase, the focus is on helping the client return to preinjury life. If that is not possible, the focus is on helping the client adjust to the changes the injury has imposed.

## Diagnostic Tests for Review

The following are routine tests done on most all hospital admissions. For this client, it is a way of monitoring the hemodynamic changes (development of anemia and so on) as well as changes in renal function. The chest x-ray lets the nurse know whether there has been an inhalation injury, a development of pneumonia, changes associated with ARDS, and so on. The complete metabolic panel gives information on electrolyte status, guiding the type of IV fluid to use, as well as whether additional electrolytes are needed. Here are the tests that should be performed:

- ▶ CBC
- ▶ Complete metabolic panel
- ▶ Urinalysis
- ▶ Chest x-ray

## Pharmacology Categories for Review

A client with burn injuries is particularly vulnerable to infection because he has lost the first line of defense, the skin. In fact, post-burn infection is a major cause of morbidity and mortality; therefore, it is helpful to review topical antibiotics used to treat those with burns. Other

complications of burns include anemia and stress ulcers. A review of medications used to treat anemia as well as medications to prevent ulcers and the bleeding that can occur will be helpful. Narcotic analgesics—particularly opiate derivatives—are used in controlling pain and providing sedation during the emergent and intermediate phases of burn care. A review of these categories, as seen in the following list, will better prepare you to care for a client with burns:

- ▶ Topical antibiotics
- ▶ Antianemics
- ▶ Antacids
- ▶ Narcotic analgesics

## Exam Prep Questions

1. The nurse is caring for a client with an electrical burn. Which structures have the greatest risk for soft tissue injury?
  - A. Fat, tendons, and bones
  - B. Skin and hair
  - C. Nerves, muscle, and blood vessels
  - D. Skin, fat, and muscle
2. Which laboratory result would be expected during the emergent phase of a burn injury?
  - A. Glucose 100 mg/dl
  - B. Potassium 3.5 mEq/l
  - C. Sodium 142 mEq/l
  - D. Albumin 4.2 gm/dl
3. An African American client is admitted with full thickness burns over 40% of his body. In addition to the CBC and complete metabolic panel, the physician is likely to request which additional bloodwork?
  - A. Erythrocyte sedimentation rate
  - B. Indirect Coombs
  - C. C reactive protein
  - D. Sickledex
4. A client weighing 76 kg is admitted at 0600 with a TBSA burn of 40%. Using the Parkland formula, the client's 24-hour intravenous fluid replacement should be:
  - A. 6,080 ml
  - B. 9,120 ml
  - C. 12,160 ml
  - D. 15,180 ml

5. On the third post-burn day, the nurse finds that the client's hourly urine output is 26 ml. The nurse should continue to assess the client and notify the doctor for an order to:
- A. Decrease the rate of the intravenous infusion.
  - B. Change the type of intravenous fluid being administered.
  - C. Change the urinary catheter.
  - D. Increase the rate of the intravenous infusion.
6. A Jewish client requires grafting to promote burn healing. Which graft is most likely to be unacceptable to the client?
- A. Isograft
  - B. Autograft
  - C. Homograft
  - D. Xenograft
7. During the rehabilitative phase, the client's burns become infected with *pseudomonas*. The topical dressing most likely to be ordered for the client is:
- A. Silver sulfadiazine (Silvadene)
  - B. Povidine (Betadine)
  - C. Mafenide acetate (Sulfamylon)
  - D. Silver nitrate
8. The CVP reading of a client with partial thickness burns is 6 mm H<sub>2</sub>O. The nurse recognizes that the client:
- A. Needs additional fluids
  - B. Has a normal CVP reading
  - C. May show signs of congestive failure
  - D. Would benefit from a diuretic
9. The physician has prescribed Protonix (pantoprazole) for a client with burns. The nurse recognizes that the medication will help prevent the development of:
- A. Curling's ulcer
  - B. Myoglobinuria
  - C. Hyperkalemia
  - D. Paralytic ileus

10. The nurse has just completed the dressing change for a client with burns to the lower legs and ankles. The nurse should place the client's ankles in which position?
- A. Internal rotation
  - B. Abduction
  - C. Dorsiflexion
  - D. Hyperextension

## Answer Rationales

1. Answer A is correct. Fat, tendon, and bone have the most resistance. The higher the resistance, the greater the heat generated by the current, thereby increasing the risk for soft tissue injury. Answer B has intermediate resistance, so it is incorrect. Answer C is incorrect because it has very low resistance. Answer D has low to intermediate resistance, so it is incorrect.
2. Answer A is correct. Glucose levels rise as a result of the stress response during the emergent phase. Answers B, C, and D are within normal range.  $K^+$  and  $Na^+$  would be elevated, whereas albumin would be lowered during the emergent period due to increased permeability.
3. Answer D is correct. Sickle cell anemia and sickle cell trait are more prevalent in African American clients. The Sickledex test detects the presence of sickle cell anemia and sickle cell trait. Trauma can trigger a sickle cell crisis, which would complicate the treatment of the client. Answers A and C indicate inflammation, so they are incorrect. Answer B is incorrect because it detects circulating antibodies against RBCs.
4. Answer C is correct. The Parkland formula is  $4 \text{ ml} \times \text{kg} \times \text{TBSA} = 24\text{-hr. fluid requirement}$ , or  $4 \times 76 \times 40 = 12,160 \text{ ml}$ . Answer A is the fluid requirement for the first 8 hours after burn injury, so it's incorrect. Answer B is incorrect because it's the fluid requirement for 16 hours after burn injury. Answer D is an excessive amount given the client's weight and TBSA, so it's incorrect.
5. Answer D is correct. The urinary output should be maintained between 30 ml and 50 ml per hour. The first action should be to increase the IV rate to prevent increased acidosis. Answer A would lead to diminished output, so it is incorrect. There is no indication that the type of IV fluid is not appropriate as is suggested by answer B, making it incorrect. Answer C would not increase the client's output and would place the client at greater risk for infection, so it is incorrect.
6. Answer D is correct. Xenografts are taken from nonhuman sources. The most common sources are porcine, or pigskin, which would be offensive to both Jews and Muslims. Answer A refers to a graft taken from an identical twin, making it incorrect. Answer B

is incorrect because it refers to a graft taken from the client's own skin. Answer C refers to a graft taken from a cadaver, making it incorrect.

7. Answer C is correct. Sulfamylon is effective in treating wounds infected with *pseudomonas*. The client should receive pain medication prior to dressing changes because the medication produces a burning sensation when applied to the wound. Answers A, B, and D are incorrect because they are used in the treatment of burns but are not effective against *pseudomonas* infections.
8. Answer B is correct. The normal CVP reading is 5–12 mm H<sub>2</sub>O. Answer A is incorrect because the client does not need additional fluids. Answers C and D would be appropriate only if the CVP reading were greater than 12 mm H<sub>2</sub>O.
9. Answer A is correct. Curling's ulcer, a stress ulcer, is a common occurrence in clients with burns. Protonix, a proton pump inhibitor, is effective in preventing ulcer formation. Answers B, C, and D are common in clients with burns but are not prevented by the use of Protonix, so they are incorrect.
10. Answer C is correct. Placing the ankles in dorsiflexed position helps prevent contractures. Answers A, B, and D will lead to contractures that may require surgical intervention, so they are incorrect.

## Suggested Reading and Resources

- ▶ Ignatavicius, D. and Workman, S. *Medical Surgical Nursing: Critical Thinking for Collaborative Care*, 5<sup>th</sup> ed. Philadelphia: Mosby, 2006.
- ▶ Brunner, L. & Suddarth, D. *Textbook of Medical Surgical Nursing*, 10<sup>th</sup> ed. Philadelphia: Lippincott Williams & Wilkins, 2006.
- ▶ Burn Recovery Center: [www.burn-recovery.org](http://www.burn-recovery.org).



# Index

## A

---

- ABCD (asymmetrical, border, colors, diameter) in assessing skin lesions, 136**
- abducens nerve assessment, 243**
- abortions, 288**
- abruptio placenta, 291**
- absence seizures, 238**
- abstinence, 304**
- acetaminophen overdose, 341**
- acid/base balance, electrolyte balance, and fluid disorders, 84**
  - changes associated with aging, 92
  - exam prep questions, 93-96
  - metabolic acidosis, 85-87
  - metabolic alkalosis, 89-90
  - normal electrolyte values, 91-92
  - pH regulation, 85
  - respiratory acidosis, 87-88
  - respiratory alkalosis, 90-91
- acidosis**
  - metabolic acidosis, 85-87
  - respiratory acidosis, 87-88
  - uncompensated acidosis, 85
- acids, 84**
- acoustic nerve assessment, 243**
- acquired heart disorders**
  - KD (Kawasaki disease), 336-338
  - rheumatic fever, 335-336
- acquired immuno-deficiency syndrome (AIDS), 294**
- acrocyanosis, 302**
- acromegaly, 200**
- active transport, 84**
- acute diarrheal disease, 334**

acute epiglottitis

**acute epiglottitis, 331-332**

**acute glaucoma, 118**

**acute glomerulonephritis, 58-59**

**acute mania, 268**

**acute otitis media (AOM), 329**

**acute post-traumatic stress disorder (PTSD), 259**

**acute respiratory failure**

ARDS (acute respiratory distress syndrome), 42

pulmonary embolus, 42-44

**acute respiratory infections**

pleurisy, 47-48

pneumonia, 46-47

TB (tuberculosis), 48

**acute subdural hematomas, 240**

**acyanotic, 325**

**Adams position, 338**

**Addiction Research Foundation Chemical Institute Withdrawal Assessment-Alcohol (CIWA-Ar), 271**

**Addisonian Crises, 206**

**ADHD (attention deficit hyperactive disorder), 276**

**administering medications, 17**

**adolescents**

growth and development, 317

psychiatric disorders, 275-276

**adrenal gland disorders, 206-207**

**adrenocortical insufficiency (Addison's disease), 206-207**

**adrenocorticotrophic hormone, 200**

**adverse effects of medications, 14**

angiotensin-converting agents, 19

anti-infectives, 21

benzodiazepines, 23

beta adrenergic blockers, 20

cholesterol-lowering agents, 28

glucocorticoids, 25-26

phenothiazines, 24

protease inhibitors, 27

proton pump inhibitors, 33

**affect, 266**

**aging clients, fluid, electrolyte balance, and acid/base balance, 92**

**AIDS (acquired immuno-deficiency syndrome), 294**

**air emboli, 43**

**alcoholism, 270-271**

**alkalosis**

metabolic alkalosis, 89-90

respiratory alkalosis, 90-91

uncompensated alkalosis, 85

**allergenic (extrinsic) asthma, 45**

**allogenic transplants, 141**

**allografts, 107**

**Allopurinol (Zyloprim), 185**

**alopecia, 139**

**alpha interferon injections, 160**

**alpha-fetoprotein screening, 285-286**

**Alzheimer's disease, 249-250**

**ambivalence, 266**

**ambulating clients, 374**

**ambulation, assistive devices for**

canes, 191

crutches, 190-191

walkers, 191-192

**American Cancer Society's seven warning signs of cancer, 134**

**American Indians. *See* Native Americans**

**aminoglycosides, 20-22**

**aminophylline, 45**

**amniocentesis, 286**

**amniotic membrane dressings (burns), 107**

**amputations, 189-190**

**analgesics, 16**

**anemia**

aplastic anemia, 73

Cooley's anemia, 75

- iron deficiency anemia, 74
- pernicious anemia, 72-73
- sickle cell anemia, 74
- aneurysm, 227-228**
- angina pectoris, 223**
- Angiotensin converting enzyme inhibitors, 219**
- angiotensin receptor blockers, 29-30, 219**
- angiotensin-converting agents, 18-19**
- anions, 84**
- anorexia nervosa, 276**
- antacids, 15**
- anti-infectives, 15, 20-22**
- antianemics, 15**
- antianxiety drugs, 22-24**
- anticholinergics, 16**
- anticoagulants, 15, 33-34, 226**
- anticonvulsants, 16, 22-23**
- antidiarrheals, 15**
- antiemetics, 24**
- antihistamines, 15**
- antihypertensives, 15, 18-19, 219**
- antipsychotics, 24, 267**
- antipyretics, 15**
- antisocial personality disorder, 264**
- anxiety-related disorders, 258**
  - DID (dissociative identity disorder), 259-260
  - GAD (generalized anxiety disorder), 258-259
  - OCD (obsessive-compulsive disorder), 261-262
  - panic disorder, 260-261
  - phobic disorders, 261
  - PTSD (post-traumatic stress disorder), 259
  - somatoform disorder, 260
- AOM (acute otitis media), 329**
- aorta, coarctation of, 326**
- APGAR scoring, 302-303**
- aplastic anemia, 73**
- Arab-Americans, 357**
  - childbirth and pain response, 358
  - dietary practices, 360
  - nonverbal/verbal communication, 358
  - time considerations, 358
- ARDS (acute respiratory distress syndrome), 42**
- arthritis, 186-187**
- artificial skin, 107**
- Asian-Americans, 355**
  - childbirth and pain response, 357
  - dietary practices, 359
  - nonverbal/verbal communication, 357
  - time considerations, 356
- assault, 372**
- assault and battery, 372**
- assistive devices for ambulation**
  - canes, 191
  - crutches, 190-191
  - walkers, 191-192
- association, 266**
- asthma, 45**
- atopic asthma, 45**
- atrioventricular node, 219**
- atropic (dry) macular degeneration, 120**
- attention deficit hyperactive disorder (ADHD), 276**
- atypical antipsychotics, 267**
- aura, 237**
- autism, 266**
- autologous transplants, 141**
- automaticisms, 238**
- autonomic hyperreflexia, 248**
- AV (atrioventricular) node, 219**
- avoidant personality disorder, 265**

## B

---

### B vitamins, 73

background diabetic retinopathy, 119

bacterial pneumonia, 46

balance suspension, 180

barbiturate abuse, 273

barrier methods of contraception, 305

bases, 84

beliefs (clients), 350

beneficial beliefs, 350

benign prostatic hyperplasia (BPH), 63

benzodiazepine abuse, 273

benzodiazepines, 22-23

beta adrenergic blockers, 19-20

beta blockers, 219

biliary atresia, 321

bilis, 353

biologic dressings, 107-108

biosynthetic dressings, 107-108

bipolar disorders

acute mania, 268

major depression, 269-270

birth control pills, 305

birth defects. *See* congenital anomalies

bladder cancer, 64-65

blood pressure, hypertension, 218-219

blood urea nitrogen (BUN), 58

blue bloaters, 44

blue spells, 326

body language

Asian-Americans, 357

Hispanics/Latinos, 353

Native Americans, 355

bone marrow transplantation, 140-142

borderline personality disorder, 264

botulism, 170

BP (blood pressure), 218-219

BPH (benign prostatic hyperplasia), 63

bradycardia, 299

brain injuries

epidural hematomas, 239

subdural hematomas, 239-240

treatment, 240

breathing, burn treatment, 103

Brethine (terbutaline sulfate), 295

bronchiolitis, 332-333

bronchitis, chronic, 44

bronchodilators, 15, 45

Buck's traction, 179

Buddhist clients, 356, 360

Buerger's disease, 226

bulimia nervosa, 276

BUN (blood urea nitrogen), 58

burns, 97-98

carbon monoxide poisoning, 103

debridement, 106

deep partial thickness (second degree), 99

diagnostic tests, 108

dressings

biologic or biosynthetic dressings, 107-108

standard wound dressings, 107

electrical burns, 99

emergent phase

additional interventions, 105

assessment, 102-103

fluid replacement formulas, 103-105

exam prep questions, 110-113

fluid replacement formulas

Consensus formula, 104-105

Parkland formula, 103-104

full thickness (third degree), 99-100

infections, 106

intermediate phase, 106-107

Lund and Browder classification method, 100

major burns, 101

minor burns, 101  
 moderate burns, 101  
 palm classification method, 101  
 pharmacology categories, 108  
 psychological care, 102  
 rehabilitative phase, 108  
 Rule of Nines, 100  
 superficial partial thickness  
 (first degree), 99  
 total body surface area (TBSA),  
 98-101

## C

### calcium

channel blockers, 219  
 normal electrolyte values, 91

### calculating fluid requirements (burns)

Consensus formula, 104-105  
 Parkland formula, 103-104

### cancer

bone marrow transplantation,  
 140-142  
 carcinoma, 134  
 chemotherapy, 137-139  
 diagnostic tests, 143-144  
 exam prep questions, 146-149  
 leukemia, 134, 340  
 lymphoma  
 definition of, 134  
 Hodgkin's lymphoma, 142-143  
 metastasis, 134  
 osteogenic sarcoma, 340  
 patient teaching, 137  
 pharmacology categories, 144-145  
 prevention, 137  
 radiation therapy, 137-138  
 risk factors, 135-136  
 sarcoma, 134  
 surgery, 137  
 TPN (total parenteral nutrition),  
 139-140

warning signs, 134  
 Wilms tumor, 340

### canes, 191

### cannabis abuse, 274-275

### caput succedaneum, 302

### carbon monoxide poisoning, 103

### carcinoma, 134

### cardiac catheterization, 225

### cardiac tamponade, 225

### cardiovascular disorders, 217-218

acquired heart disorders  
 KD (Kawasaki disease), 336-338  
 rheumatic fever, 335-336  
 aneurysm, 227-228  
 Buerger's disease, 226  
 CHD (congenital heart defects),  
 324-326  
 acyanotic, 325  
 COA (coarctation of the aorta),  
 326  
 cyanotic, 325  
 symptoms, 325  
 TOF (tetralogy of Fallot),  
 326-327  
 congestive heart failure, 228-229  
 diagnostic tests, 229  
 exam prep questions, 230-232  
 heart block  
 first-degree, 219  
 pacemakers/internal defibrillators,  
 221-222  
 second-degree, 220  
 third-degree, 220  
 toxicity to medications, 221  
 hypertension, 218-219  
 myocardial infarction, 222  
 angina pectoris, 223  
 diagnosis, 223  
 managing, 224  
 ventricular fibrillation (V-fib),  
 224-226  
 ventricular tachycardia, 224

## cardiovascular disorders

- pharmacology categories, 229
  - Raynaud's Syndrome, 227
  - thrombophlebitis, 227
- carditis, 336**
- casts, 181**
- CAT (Computer Adaptive Test), 4-5**
- cataracts, 116-117**
- catatonic schizophrenia, 267**
- Category A (pregnancy drug category), 36**
- Category B (pregnancy drug category), 36**
- Category C (pregnancy drug category), 36**
- Category D (pregnancy drug category), 36**
- Category X (pregnancy drug category), 36**
- Catholic clients, 360**
- cations, 84**
- celiac, 335**
- central nervous system disorders, congenital anomalies, 323**
- central venous pressure (CVP), 106**
- cephalocaudal – proximodistal development, 312**
- cephalohematoma, 302**
- cerebral perfusion pressure (CPP), 245**
- cervical mucus method of contraception, 304**
- CHD (congenital heart defects), 324**
  - acyanotic, 325
  - coarctation of the aorta (COA), 326
  - cyanotic, 325
  - symptoms, 325
  - tetralogy of Fallot (TOF), 326-327
- Chemical Institute Withdrawal Assessment-Alcohol (CIWA-Ar), 271**
- chemical names, 18**
- chemotherapy, 137, 139**
- childbirth. *See also* pregnancy**
  - Arab-Americans, 358
  - Asian-Americans, 357
  - Hispanics/Latinos, 354
  - labor
    - definition of, 295
    - dilation, 297
    - dystocia, 297
    - effacement, 297
    - factors influencing, 295-296
    - fetal lie, 297
    - fetal monitoring, 298-300
    - pharmacologic management, 301-302
    - phases, 296
    - position, 296
    - precipitate delivery, 297
    - prelabor testing, 298
    - presentation, 296
    - preterm labor, 295
    - stages, 296
    - station, 297
  - Native Americans, 355
  - postpartum care, 302
  - terms associated with newborns, 302-303
- childhood psychiatric disorders, 275**
  - ADHD (attention deficit hyperactive disorder), 276
  - conduct disorder, 275
  - eating disorders, 276
  - oppositional defiant disorder, 275
- children. *See* pediatric clients**
- Chinese. *See* Asian-Americans**
- chlamydia, 292**
- chloride, normal electrolyte values, 91**
- cholecystitis, 167-168**
- cholelithiasis, 167-169**
- cholesterol-lowering agents, 28-29**
- chronic bronchitis, 44**
- chronic glomerulonephritis, 59**
- chronic obstructive pulmonary disease (COPD)**
  - asthma, 45
  - chronic bronchitis, 44
  - emphysema, 44-45

- chronic subdural hematomas, 240**
- Chvostek's sign, 90, 204**
- circulation, burn treatment, 103**
- cirrhosis, 163-165**
- civil laws, 369**
- CIWA-Ar (Chemical Institute Withdrawal Assessment-Alcohol), 271**
- classifications of burns, 98**
  - deep partial thickness (second degree), 99
  - full thickness (third degree), 99
  - Lund and Browder method, 100
  - palm method, 101
  - Rule of Nines, 100
  - superficial partial thickness (first degree), 99
- cleft lip, 318-319**
- cleft palate, 318-319**
- client beliefs, 350**
- client care, managing, 373-376**
- client needs exam prep questions, 9-12**
- client rights, 370**
- closed fractures, 178**
- clubfoot, congenital, 323**
- Cluster A personality disorders, 262-263**
- Cluster B personality disorders, 263-264**
- Cluster C personality disorders, 265**
- CNS. *See* central nervous system**
- COA (coarctation of the aorta), 326**
- Code of Ethics for Nursing, 370-371**
- codeine abuse, 273**
- coining, 356**
- coitus interruptus, 304**
- comminuted fractures, 178**
- common laws, 369-370**
- communication**
  - Arab-Americans, 358
  - Asian-Americans, 357
  - Hispanics/Latinos, 353
  - language differences, 351
  - Native Americans, 355
- compartment syndrome, 181-182**
- complete abortions, 288**
- complete spinal cord injuries, 246**
- complex partial seizures, 238**
- complications of pregnancy, 287**
- compound fractures, 178**
- Computer Adaptive Test (CAT), 4-5**
- condoms, 305**
- conduct disorder, 275**
- condylomata, 293**
- congenital aganglionic megacolon (Hirschsprung disease), 320-321**
- congenital anomalies, 317-318**
  - biliary atresia, 321
  - cleft lip and cleft palate, 318-319
  - congenital clubfoot, 323
  - CHD (congenital heart defects), 324-326
    - acyanotic, 325
    - COA (coarctation of the aorta), 326
    - cyanotic, 325
    - symptoms, 325
    - TOF (tetralogy of Fallot), 326-327
  - DHD (developmental hip dysplasia), 322
  - EA (esophageal atresia), 319
  - galactosemia, 328
  - Hirschsprung disease (congenital aganglionic megacolon), 320-321
  - imperforate anus, 320
  - PKU (phenylketonuria), 327-328
  - spina bifida, 323-324
  - TEF (tracheoesophageal fistula), 319
- congestive heart failure, 228-229**
- Consensus formula, 104-105**
- consent for care, witnessing, 373**
- continuous passive motion (CPM) machine, 188-189**
- contraception, 304-305**

contraction stress test

**contraction stress test, 298**  
**contusions of eyes, 122**  
**Cooley's anemia, 75**  
**COPD (chronic obstructive pulmonary disease)**  
     asthma, 45  
     chronic bronchitis, 44  
     emphysema, 44-45  
**cord prolapse, 291**  
**coronary artery bypass grafts, 225**  
**cortisol, 206**  
**Cotrel-Duboussset approach, 339**  
**Coumadin, 226**  
**cox 2 enzyme blockers, 30-31**  
**coxa plana, 339**  
**CPM (continuous passive motion machine, 188-189)**  
**CPP (cerebral perfusion pressure), 245**  
**cranial nerve assessment, 243-244**  
**craniotomy (intracranial surgery), 245-246**  
**cretinism, 201-202**  
**criminal laws, 369**  
**Crohn's disease, 155**  
**crutches, 190-191**  
**Crutchfield tong traction, 180**  
**cultural practices**  
     Arab-Americans, 357-358  
     Asian-Americans, 355-357  
     cultural assessment  
         client beliefs, 350  
         language differences, 351  
     dietary considerations, 359-360  
     exam prep questions, 362-365  
     Hispanics/Latinos, 352-354  
     Native Americans, 354-355  
     religious beliefs, 360-361  
**cultured skin dressings, 107**  
**cupping, 356**  
**Cushing's Syndrome, 26**

**Cushing's triad, 241**  
**CVP (central venous pressure), 106**  
**cyanotic, 325**  
**Cyclogyl, 117**  
**cystectomy, 64**  
**cystic fibrosis, 333**

## D

---

**debridement, 106**  
**deceleration of fetal heart tones, 299-300**  
**deep partial thickness (second degree) burns, 99**  
**defibrillators, internal, 221-222**  
**degenerative neurological disorder, 249-250**  
**delayed post-traumatic stress disorder (PTSD), 259**  
**demand pacemakers, 221**  
**Demerol, 14**  
**dependent personality disorder, 265**  
**depression, 269-270**  
**determiners, 7**  
**development and growth**  
     adolescents, 317  
     infants, 312-313  
     preschoolers, 315-316  
     school age children, 316  
     toddlers, 314  
**dextrostix, 211**  
**DHD (developmental hip dysplasia), 322**  
**diabetes during pregnancy, 289**  
**diabetes mellitus, 207, 209-210**  
**diabetic retinopathy, 119-120**  
**Diagnostic and Statistical Manual of Mental Disorders (DSM-IV-TR), 258**  
**diagnostic tests**  
     burns, 108  
     cancer, 143-144  
     cardiovascular disorders, 229  
     ear disorders, 126



endocrine system disorders, 210-211  
 gastrointestinal disorders, 170-171  
 hematopoietic disorders, 76  
 musculoskeletal disorders, 192-193  
 neurological disorders, 250-251  
 obstetric clients, 306  
 pediatric clients, 342  
 prenatal care, 286  
 psychiatric disorders, 277  
 renal/genitourinary disorders, 65  
 respiratory disorders, 50  
 ulcers, 153  
 visual tests, 123

**dialysis, 60**

**Diamox, 117**

**diaphragms, 305**

**diastolic pressure, 218**

**DIC (disseminated intravascular coagulation), 290**

**DID (dissociative identity disorder), 259-260**

**diet. *See* nutrition**

**dietary practices of cultural groups, 359-360**

**dilation, 297**

**dissecting aneurysm, 227**

**disseminated intravascular coagulation (DIC), 290**

**dissociative identity disorder (DID), 259-260**

**distractors, 7**

**diuretics, 15, 219**

**diversity. *See* cultural practices**

**diverticulitis, 157**

**dressings**

- burn dressings, 107-108
- TPN (total parenteral nutrition), 140

**drugs. *See* pharmacology**

**dry (atropic) macular degeneration, 120**

**DSM-IV-TR (Diagnostic and Statistical Manual of Mental Disorders), 258**

**dumping syndrome, 154**

**duodenal ulcers, 152**

**dysreflexia, 248**

**dysrhythmias, 224-226**

**dystocia, 297**

## E

---

**e. coli, 170**

**EA (esophageal atresia), 319**

**ear disorders, 123**

- diagnostic tests, 126
- ear trauma, 126
- exam prep questions, 128-130
- hearing loss, 126
- Meniere's disease, 124-125
- otitis externa, 124
- otitis media, 124
- otosclerosis, 125
- pharmacology categories, 127
- presbycusis, 125

**early deceleration of fetal heart tones, 299**

**eating disorders, 276**

**ECCE (extracellular cataract extraction), 117**

**echinacea, 35**

**effacement, 297**

**elective abortions, 288**

**electrical burns, 99**

**electrocardiograms, 220-221**

**electrolyte balance, acid/base balance, and fluid disorders, 84**

- changes associated with aging, 92
- exam prep questions, 93-96
- metabolic acidosis
  - care and treatment, 86-87
  - causes, 85-86
  - definition of, 85
  - symptoms, 86
- metabolic alkalosis, 89-90

electrolyte balance, acid/base balance, and fluid disorders

normal electrolyte values, 91-92

pH regulation, 85

respiratory acidosis, 87-88

respiratory alkalosis, 90-91

### **emergent phase (burns)**

additional interventions, 105

assessment, 102-103

fluid replacement formulas, 103-105

### **emerging infections, 48-49**

### **empacho, 353**

### **emphysema, 44-45**

### **end stage renal disease (ESRD), 60-61**

### **endocrine system disorders, 199**

adrenal gland disorders, 206-207

diabetes mellitus, 207, 209-210

diagnostic tests, 210-211

exam prep questions, 212-215

parathyroid disorders, 204-205

pharmacology categories, 211

pituitary disorders, 200-201

thyroid disorders, 201-203

### **engrafts, 141**

### **enteric-coated tablets, 16**

### **epidural block, 301**

### **epidural hematomas, 239**

### **epiglottitis, acute, 331-332**

### **erythema marginatum, 336**

### **erythroblastosis fetalis, 303**

### **esophageal atresia (EA), 319**

### **ESRD (end stage renal disease), 60**

hemodialysis, 60

peritoneal dialysis, 60

renal transplants, 61

### **ethics, Code of Ethics for Nursing, 370-371**

### **evil eye (mal de ojo), 353**

### **exam. See NCLEX-PN exam**

### **extracellular cataract extraction (ECCE), 117**

### **extracellular fluid, 84**

### **extrinsic (allergenic) asthma, 45**

### **exudative (wet) macular degeneration, 120**

### **eye disorders, 116**

exam prep questions, 128-130

intraocular disorders

cataracts, 116-117

glaucoma, 117-119

pharmacology categories, 123

refractory errors, 121-122

retinal disorders, 119-120

traumatic injuries, 122

visual tests, 123

## **F**

### **facial nerve assessment, 243**

### **farsightedness, 121**

### **fasciotomy, 182**

### **fasting blood glucose, 211**

### **fat emboli, 42-43**

### **felonies, 369**

### **fetal heart tones, measuring, 287**

### **fetal lie, 297**

### **fetal monitoring, 298-300**

### **feverfew, 35**

### **filtration, 84**

### **first degree (superficial partial thickness) burns, 99**

### **first-degree heart block, 219**

### **fluid replacement formulas (burns)**

Consensus formula, 104-105

Parkland formula, 103-104

### **fluid, electrolyte balance, and acid/base balance disorders, 84**

changes associated with aging, 92

exam prep questions, 93-96

metabolic acidosis

care and treatment, 86-87

causes, 85-86

- definition of, 85
- symptoms, 86
- metabolic alkalosis, 89-90
- normal electrolyte values, 91-92
- pH regulation, 85
- respiratory acidosis, 87-88
- respiratory alkalosis, 90-91
- focal seizures, 238**
- food-borne illnesses, 169-170**
- foreign bodies in eye, 122**
- four-point crutch-walking gait, 191**
- fractured hip, 187-188**
- fractures**
  - casts, 181
  - comminuted fractures, 178
  - compartment syndrome, 181-182
  - compound fractures, 178
  - definition of, 178
  - green stick fractures, 178
  - osteomyelitis, 182-183
  - pathological fractures, 178
  - simple fractures, 178
  - symptoms, 178
  - traction, 179-180
  - treatment, 178-179
- fraud, 373**
- full thickness (third degree) burns, 99-100**
- fusiform aneurysm, 227**

## G

---

- GAD (generalized anxiety disorder), 258-259**
- gait belts, 192**
- galactosemia, 328**
- gallbladder disease, 167**
- gallbladder inflammation, 167-168**
- garamycin, 47**
- gastric ulcers, 153**
- gastroenteritis, 334**

## **gastrointestinal disorders**

- celiac, 335
- cholecystitis, 167-168
- cholelithiasis, 167-169
- cirrhosis, 163-165
- congenital anomalies
  - biliary atresia, 321
  - cleft lip and cleft palate, 318-319
  - EA (esophageal atresia), 319
  - Hirschsprung disease (congenital aganglionic megacolon), 320-321
  - imperforate anus, 320
  - TEF (tracheoesophageal fistula), 319
- Crohn's disease, 155
- diagnostic tests, 170-171
- diverticulitis, 157
- exam prep questions, 172-175
- food-borne illnesses, 169-170
- gastroenteritis, 334
- hepatitis
  - general management techniques, 158
  - hepatitis A, 158-159
  - hepatitis B, 159-161
  - hepatitis C, 161
  - hepatitis D, 161-162
  - hepatitis E, 162
  - icteric stage, 162-163
  - prodromal stage, 162-163
- intussusception, 334
- pancreatitis, 165-166
- pharmacology categories, 171
- pyloric stenosis, 334
- ulcerative colitis, 156
- ulcers
  - diagnostic tools, 153
  - dumping syndrome, 154
  - duodenal ulcers, 152
  - gastric ulcers, 153
  - treatment, 153-154

general anesthesia, childbirth

**general anesthesia, childbirth, 302**

**generalized anxiety disorder (GAD), 258-259**

**generalized seizures**

absence, 238

tonic-clonic, 236-237

**generic names, 18**

**genital herpes, 293**

**genitourinary disorders, 57**

acute glomerulonephritis, 58-59

bladder cancer, 64-65

BPH (benign prostatic hyperplasia), 63

chronic glomerulonephritis, 59

diagnostic tests, 65

ESRD (end stage renal disease), 60-61

exam prep questions, 66-69

nephrotic syndrome, 61-62

pharmacology categories, 65

urinary calculi, 62

UTI (urinary tract infections), 62-63

**gigantism, 200**

**ginko, 35**

**ginseng, 35**

**Glasgow coma scale, 244-245**

**glaucoma**

acute glaucoma, 118

management of, 118-119

POAG (primary open-angle glaucoma), 117

secondary glaucoma, 118

**glomerulonephritis, 58-59**

**glossopharyngeal nerve assessment, 243**

**glucagon, 210**

**glucocorticoids, 25**

**glucose tolerance tests, 210**

**gluten-induced enteropathy, 335**

**glycosylated hemoglobin (Hgb A-1C), 211**

**goiters, 202**

**gonadotrophic hormones, 200**

**gonorrhea, 292**

**gout, 184-185**

**Gower's maneuver, 339**

**grand mal seizures, 236-237**

**green stick fractures, 178**

**growth and development**

adolescents, 317

infants, 312-313

preschoolers, 315-316

school age children, 316

toddlers, 314

**growth hormone, 200**

**Guillain-Barre, 248-249**

**Guthrie test, 328**

## H

**H Pylori bacteria, 152**

**H.influenza B conjugate vaccine, 331**

**hallucinogens abuse, 274**

**halo vests, 247**

**Harrington rods, 339**

**Havrix, 159**

**hazardous substances, ingestion of**

acetaminophen overdose, 341

iron poisoning, 342

lead, 341

salicylate overdose, 341

**HBIG (hepatitis B immune globulin), 161**

**hearing loss, assisting clients with, 126.**

**See also ear disorders**

**heart block**

first-degree, 219

pacemakers/internal defibrillators, 221-222

second-degree, 220

third-degree, 220

toxicity to medications, 221

**HELLP syndrome, 290**

**hematomas**

- epidural hematomas, 239
- subdural hematomas, 239-240
- treatment, 240

**hematopoietic disorders, 71**

- anemia, 72
  - aplastic anemia, 73
  - Cooley's anemia, 75
  - iron deficiency anemia, 74
  - pernicious anemia, 72-73
  - sickle cell anemia, 74
- diagnostic tests, 76
- exam prep questions, 79-81
- hemophilia, 75
- pharmacology categories, 77
- polycythemia vera, 76

**hemodialysis, 60****hemolysis, 290****hemophilia, 75****heparin, 226****hepatitis**

- general management techniques, 158
- hepatitis A, 158-159
- hepatitis B, 159-161
- hepatitis C, 161
- hepatitis D, 161-162
- hepatitis E, 162
- icteric stage, 162-163
- prodromal stage, 162-163

**hepatitis B immune globulin (HBIG), 161****Heptovax, 160****herbals, 35-36****herpes, 293****heterograftsw, 107****HEV (hepatitis E), 162****Hgb A-1C (glycosylated hemoglobin), 211****high-purine foods, 185****Hindu clients, 356, 361****hip fractures, 187-188****hip replacement, 187-188****Hirschsprung disease (congenital aganglionic megacolon), 320-321****Hispanics/Latinos**

- bilis, 353
- childbirth and pain response, 354
- dietary practices, 359
- empacho, 353
- evil eye (mal de ojo), 353
- nonverbal/verbal communication, 353
- susto (fright sickness), 353
- time considerations, 353
- traditional healers, 352

**histamine 2 antagonists, 31-32****histrionic personality disorder, 263****HIV (human immunodeficiency virus), 294****Hodgkin's lymphoma, 142-143****hormonal contraception, 305****hormones**

- adrenocorticotrophic hormone, 200
- cortisol, 206
- gonadotrophic hormone, 200
- growth hormone, 200
- parathormone, 204-205
- thyroid hormone, 201
- thyrotrophic hormone, 200

**human immunodeficiency virus (HIV), 294****hydatidiform moles, 288****hyperbilirubinemia, 303****hyperemesis gravidarum, 287****hyperglycemia, 209****hyperkalemia, 59, 86****hyperopia, 121****hyperparathyroidism, 204-205****hyperplasia of the thyroid, 202****hypertension, 218-219****hypertensive retinopathy, 119****hyperthyroidism, 202-203****hyphema, 122****hypoglossal nerve assessment, 244**

hypoglycemia

**hypoglycemia, 210****hypoparathyroidism, 204-205****hypothyroidism, 201-202****I****ICP (increased intracranial pressure)**

causes, 240

intracranial pressure monitors, 245

symptoms, 240-242

treatment, 242

**icteric stage (hepatitis), 162-163****identifying drug types, 18, 34-35****Imferon, 75****imperforate anus, 320****incompetent cervix, 287****incomplete abortions, 288****incomplete spinal cord injuries, 246****inevitable abortions, 288****infants. *See also* obstetric clients, pregnancy; pediatric clients**

acrocyanosis, 302

APGAR scoring, 302-303

caput succedaneum, 302

cephalohematoma, 302

growth and development, 312-313

hyperbilirubinemia, 303

ICP (increased intracranial pressure), 242

milia, 303

Mongolian spots, 303

physiologic jaundice, 304

prematurity, 295

**infection control, 8****infections during pregnancy, 291-294****inflammatory bowel disorders, 154**

Crohn's disease, 155

diverticulitis, 157

ulcerative colitis, 156

**ingestion of hazardous substances**

acetaminophen overdose, 341

iron poisoning, 342

lead, 341

salicylate overdose, 341

**insulin, 209****intact corneal rings, 122****Integra, 107****intermediate phase (burns), 106-107****internal defibrillators, 221-222****intracellular fluid, 84****intracranial pressure monitors, 245****intracranial surgery (craniotomy), 245-246****intramuscular iron, 75****intraocular disorders**

cataracts, 116-117

glaucoma

acute glaucoma, 118

management of, 118-119

POAG (primary open-angle glaucoma), 117

secondary glaucoma, 118

**intraocular pressure, 118****intrapartal care**

dilation, 297

dystocia, 297

effacement, 297

fetal lie, 297

fetal monitoring, 298-300

labor

definition of, 295

factors influencing, 295-296

pharmacologic management, 301-302

phases, 296

stages, 296

position, 296

precipitate delivery, 297

prelabor testing, 298

presentation, 296

station, 297

intrauterine devices (IUDs), 305  
 intrinsic (nonallergenic) asthma, 45  
 intussusception, 334  
 iron deficiency anemia, 74  
 iron poisoning, 342  
 Islamic clients, 357  
 IUD (intrauterine devices), 305  
 IV therapy, 374

## J - K

---

jabon de la mano milagrosa, 353  
 Japanese. *See* Asian-Americans  
 jaundice, 304, 322  
 Jehovah's Witness clients, 361  
 Jewish clients, 361

kava-kava, 36  
 KD (Kawasaki disease), 336-338  
 kernicterus, 303  
 ketonuria, 208  
 keywords, looking for, 6  
 kidney stones, 62. *See also* renal/genitourinary disorders  
 knee replacement, 188-189

## L

---

lacerations of eye, 122  
 language differences, 351  
 Lantus insulin, 209  
 laryngotracheobronchitis (LTB), 331  
 LASIK (laser in-situ keratomileusis), 122  
 Latinos/Hispanics
 

- bilis, 353
- childbirth and pain response, 354
- empacho, 353
- evil eye (mal de ojo), 353
- nonverbal/verbal communication, 353

susto (fright sickness), 353  
 time considerations, 353  
 traditional healers, 352

### laws

civil laws, 369  
 common laws, 369-370  
 criminal laws, 369  
 statutory laws/regulatory laws, 369

### laxatives, 15

### lead poisoning, 341

### left occiput anterior (LOA), 296

### legal issues

assault, 372  
 assault and battery, 372  
 civil laws, 369  
 Code of Ethics for Nursing, 370-371  
 common laws, 369-370  
 criminal laws, 369  
 exam prep questions, 377-380  
 fraud, 373  
 malpractice, 372  
 managing client care, 373-376  
 negligence, 372  
 Nurse Practice Acts, 368-369  
 statutory laws/regulatory laws, 369  
 torts, 372  
 witnessing consent for care, 373

### Legg-Calve-Perthes disease, 339

### Legionnaire's Disease, 49

### Leopold's maneuver, 297

### leukemia, 134, 340

### lithium, 269

### liver disorders

cirrhosis, 163
 

- biliary cirrhosis, 163
- diagnosis, 164
- Laennec's portal cirrhosis, 163
- post-necrotic cirrhosis, 163
- symptoms, 163-164
- treatment, 164-165

## liver disorders

- hepatitis
  - general management techniques, 158
  - hepatitis A, 158-159
  - hepatitis B, 159-161
  - hepatitis C, 161
  - hepatitis D, 161-162
  - hepatitis E, 162
  - icteric stage, 162-163
  - prodromal stage, 162-163

**LOA (left occiput anterior), 296**

**local infiltration, 301**

**low-purine foods, 185**

**LTB (laryngotracheobronchitis), 331**

**Lugol's solution, 203**

**Lund and Browder classification method (burns), 100**

**lungs. *See* respiratory disorders**

**Luque wires, 339**

**lymphoma**

- definition of, 134
- Hodgkin's lymphoma, 142-143

## M

**ma huang, 36**

**macular degeneration, 120**

**magnesium, normal electrolyte values, 92**

**magnesium gluconate, 290**

**magnesium sulfate, 290, 295**

**major burns, 100-101**

- emergent phase
  - additional interventions, 105
  - assessment, 102-103
  - fluid replacement formulas, 103-105
- fluid replacement formulas
  - Consensus formula, 104-105
  - Parkland formula, 103-104

**major depression, 269-270**

**maladaptive beliefs, 350**

**malignant cells. *See* cancer**

**malpractice, 372**

**managing client care, 373-376**

**mania, acute, 268**

**Mantoux skin tests, 48**

**manual traction, 179**

**MAOI (monoamine oxidase inhibitors), 270**

**MAP (mean arterial pressure), 245**

**marijuana abuse, 274-275**

**maternal/infant clients, 283-284**

- abortions, 288
- abruptio placenta, 291
- complications of pregnancy, 287
- contraception, 304-305
- cord prolapse, 291
- diagnostic tests, 306
- disseminated intravascular coagulation (DIC), 290
- exam prep questions, 307-310
- labor
  - definition of, 295
  - dilation, 297
  - dystocia, 297
  - effacement, 297
  - factors influencing, 295-296
  - fetal lie, 297
  - fetal monitoring, 298-300
  - pharmacologic management, 301-302
  - phases, 296
  - position, 296
  - precipitate delivery, 297
  - prelabor testing, 298
  - presentation, 296
  - preterm labor, 295
  - stages, 296
  - station, 297
- maternal diabetes, 289
- maternal infections, 291-294



- pharmacological categories, 306
- physiologic jaundice, 304
- Placenta Previa, 291
- postpartum care, 302
- preeclampsia, 289-290
- prematurity, 295
- prenatal care
  - alpha-fetoprotein screening, 285-286
  - amniocentesis, 286
  - diagnostic tests, 286
  - diet and weight maintenance, 285
  - fetal heart tones, measuring, 287
  - ultrasonography, 287
- Rh incompatibility, 303
- signs of pregnancy
  - positive signs, 285
  - presumptive signs, 284
  - probable signs, 284-285
- terms associated with newborns, 302-303
- mean arterial pressure (MAP), 245**
- medication. *See* pharmacology**
- Meniere's disease, 124-125**
- meningitis, 324**
- meningocele spina bifida, 323**
- mental disorders. *See* psychiatric disorders**
- metabolic acidosis**
  - care and treatment, 86-87
  - causes, 85-86
  - definition of, 85
  - symptoms, 86
- metabolic alkalosis, 89-90**
- metabolic disorders, 327-328**
- metastasis, 134**
- methicillin-resistant staphylococcus aureus (MRSA), 22**
- milia, 303**
- minor burns, 101**
- miotics, 16**
- misdeemeanors, 369**
- missed abortions, 288**
- moderate burns, 101**
- Mongolian spots, 303**
- monoamine oxidase inhibitors (MAOI), 270**
- morphine abuse, 273**
- MRSA (methicillin-resistant staphylococcus aureus), 22**
- mucocutaneous lymph node syndrome, 336-338**
- mucoviscidosis (cystic fibrosis), 333**
- multiple personality disorder, 259-260**
- multiple sclerosis, 249-250**
- muscular dystrophies, 339**
- musculoskeletal disorders, 177, 338**
  - amputations, 189-190
  - assistive devices for ambulation
    - canes, 191
    - crutches, 190-191
    - walkers, 191-192
  - congenital anomalies
    - congenital clubfoot, 323
    - developmental hip dysplasia (DHD), 322
  - diagnostic tests, 192-193
  - exam prep questions, 194-197
  - fractures
    - casts, 181
    - comminuted fractures, 178
    - compartment syndrome, 181-182
    - compound fractures, 178
    - definition of, 178
    - green stick fractures, 178
    - osteomyelitis, 182-183
    - pathological fractures, 178
    - simple fractures, 178
    - symptoms, 178
    - traction, 179-180
    - treatment, 178-179

## musculoskeletal disorders

- gout, 184-185
- Legg-Calve-Perthes disease, 339
- muscular dystrophies, 339
- osteoporosis, 183-184
- pharmacology, 193
- rheumatoid arthritis, 186-187
- scoliosis, 338-339
- surgical procedures
  - amputations, 189-190
  - fractured hip and hip replacement, 187-188
  - total knee replacement, 188-189

**Muslim clients, 357****myasthenia gravis, 249-250****mydriatics, 16****myelomeningocele spina bifida, 323****myocardial infarction, 222**

- angina pectoris, 223
- diagnosis, 223
- managing, 224
- ventricular fibrillation (V-fib), 224-226
- ventricular tachycardia, 224

**myopia, 121****myxedema, 201**


---

## N

**narcissistic personality disorder, 263****narcotics, 16****narrow-angle glaucoma, 118****National Council Licensure Examination.***See* NCLEX-PN exam**Native Americans**

- childbirth and pain response, 355
- nonverbal/verbal communication, 355
- shamans, 354
- time considerations, 354

**NCLEX exam**

- exam prep questions
  - burns, 110-113
  - cancer, 146-149

## cardiovascular disorders, 230-232

## cultural practices, 362-365

## endocrine system disorders, 212-215

## fluid, electrolyte balance, and acid/base balance, 93-96

## gastrointestinal disorders, 172-175

## hematopoietic disorders, 79-81

## legal issues, 377-380

## musculoskeletal disorders, 194-197

## neurological disorders, 252-255

## nursing process/client needs, 9-12

## obstetric clients, 307-310

## pediatric clients, 344-346

## pharmacology, 37-40

## psychiatric disorders, 278-281

## renal/genitourinary disorders, 66-69

## respiratory disorders, 52-55

## sensorineural disorders, 128-130

## preparing for, CAT (Computer Adaptive Test), 4-5

## test-taking strategies, 5-8

**nearsightedness, 121****negative schizophrenia, 266****negligence, 372****Neo-Synephrine, 117****nephroblastoma, 340****nephrotic syndrome, 61-62****nerve blocks, 301****neuroleptic malignant syndrome, 268****neurological assessment**

- cranial nerve assessment, 243-244
- Glasgow coma scale, 244-245
- intracranial pressure monitors, 245

**neurological disorders, 235**

- brain injuries, 239-240
- degenerative neurological disorders, 249-250

- diagnostic tests, 250-251
  - exam prep questions, 252-255
  - Guillain-Barre, 248-249
  - ICP (increased intracranial pressure)
    - causes, 240
    - intracranial pressure monitors, 245
    - symptoms, 240-242
    - treatment, 242
  - intracranial surgery (craniotomy), 245-246
  - neurological assessment
    - cranial nerve assessment, 243-244
    - Glasgow coma scale, 244-245
    - intracranial pressure monitors, 245
  - pharmacology, 251
  - seizures, 236
    - absence, 238
    - tonic-clonic, 236-238
    - complex partial, 238
    - simple partial, 238
  - spinal cord injuries
    - autonomic hyperreflexia, 248
    - complete, 246
    - incomplete, 246
    - spinal shock, 248
    - treatment, 247
  - status epilepticus, 239
  - neurotic disorders. *See* anxiety-related disorders**
  - neurotransmitters, 258**
  - neutral beliefs, 350**
  - newborns. *See* infants**
  - nitroglycerine, 223**
  - non-cardiogenic pulmonary edema, 42**
  - non-stress test, 298**
  - nonallergenic (intrinsic) asthma, 45**
  - nonsteroidal anti-inflammatory drugs (NSAIDs), 30**
  - nonverbal/verbal communication**
    - Arab-Americans, 358
    - Asian-Americans, 357
    - Hispanics/Latinos, 353
    - Native Americans, 355
  - normal electrolyte values, 91-92**
  - Norplant, 305**
  - novel antipsychotics, 267**
  - NSAIDs (nonsteroidal anti-inflammatory drugs), 30**
  - Nurse Practice Acts, 369, 375**
  - Nursing Practice Acts, 368**
  - nursing process exam prep questions, 9-12**
  - nutrition**
    - prenatal diet and weight maintenance, 285
    - TPN (total parenteral nutrition), 139-140
- 
- O
- OA (occiput anterior), 297**
  - obsessive-compulsive personality disorder, 265**
  - obstetric clients, 283-284**
    - abortions, 288
    - abruptio placenta, 291
    - complications of pregnancy, 287
    - contraception, 304-305
    - cord prolapse, 291
    - diagnostic tests, 306
    - disseminated intravascular coagulation (DIC), 290
    - exam prep questions, 307-310
    - labor
      - definition of, 295
      - dilation, 297
      - dystocia, 297
      - effacement, 297
      - factors influencing, 295-296
      - fetal lie, 297
      - fetal monitoring, 298-300
      - pharmacologic management,

## obstetric clients

- 301-302
- phases, 296
- position, 296
- precipitate delivery, 297
- prelabor testing, 298
- presentation, 296
- preterm labor, 295
- stages, 296
- station, 297
- maternal diabetes, 289
- maternal infections, 291-294
- pharmacological categories, 306
- physiologic jaundice, 304
- Placenta Previa, 291
- postpartum care, 302
- preeclampsia, 289-290
- prematurity, 295
- prenatal care
  - alpha-fetoprotein screening, 285-286
  - amniocentesis, 286
  - diagnostic tests, 286
  - diet and weight maintenance, 285
  - fetal heart tones, measuring, 287
  - ultrasonography, 287
- Rh incompatibility, 303
- signs of pregnancy
  - positive signs, 285
  - presumptive signs, 284
  - probable signs, 284-285
- terms associated with newborns, 302-303
- OCD (obsessive-compulsive disorder), 261-262**
- oculomotor nerve assessment, 243**
- olfactory nerve assessment, 243**
- opiates, 273**
- oppositional defiant disorder, 275**
- optic nerve assessment, 243**
- osmosis, 84**
- osteogenic sarcoma, 340**

**osteomyelitis, 182-183****osteoporosis, 183-184****osteosarcoma, 340****otitis externa, 124****otitis media, 124****otosclerosis, 125****overdoses**

acetaminophen overdose, 341

salicylate overdose, 341

---

**P**
**pacemakers, 221-222****pain response**

Arab-Americans, 358

Asian-Americans, 357

Hispanics/Latinos, 354

Native Americans, 355

**palm classification method (burns), 101****pancreatitis, 165-166****panic disorder, 260-261****paranoid personality disorder, 262****parathormone, 204-205****parathyroid disorders, 204-205****Parkinson's disease, 249-250****Parkland formula, 103-104****partial seizures, 238****partial thromoplastin time (PTT), 226****pathological fractures, 178****Patient's Bill of Rights, 370****Pearson attachments, 180****pediatric clients**

acquired heart disorders

KD (Kawasaki disease), 336-338

rheumatic fever, 335-336

childhood cancer, 340

congenital anomalies, 317

biliary atresia, 321

cleft lip and cleft palate, 318-319

congenital clubfoot, 323

- CHD (congenital heart defects), 324-327
- DHD (developmental hip dysplasia), 322
- EA (esophageal atresia), 319
- galactosemia, 328
- Hirschsprung disease (congenital aganglionic megacolon), 320-321
- imperforate anus, 320
- PKU (phenylketonuria), 327-328
- spina bifida, 323-324
- TEF (tracheoesophageal fistula), 319
- diagnostic tests, 342
- exam prep questions, 344-346
- gastrointestinal disorders, 334-335
- growth and development
  - adolescents, 317
  - infants, 312-313
  - preschoolers, 315-316
  - school age children, 316
  - toddlers, 314
- ingestion of hazardous substances, 341-342
- musculoskeletal disorders, 338-339
- pharmacology categories, 343
- respiratory disorders
  - acute epiglottitis, 331-332
  - AOM (acute otitis media), 329
  - bronchiolitis, 332-333
  - cystic fibrosis, 333
  - LTB (laryngotracheobronchitis), 331
  - tonsillitis, 330
- penetrating injuries of eye, 122**
- peritoneal dialysis, 60**
- peritonitis, 60**
- pernicious anemia, 72-73**
- personality disorders**
  - antisocial personality disorder, 264
  - avoidant personality disorder, 265
  - borderline personality disorder, 264
  - client management, 265
  - dependent personality disorder, 265
  - histrionic personality disorder, 263
  - narcissistic personality disorder, 263
  - obsessive-compulsive personality disorder, 265
  - paranoid personality disorder, 262
  - schizoid personality disorder, 263
  - schizotypal personality disorder, 263
- PE (polyethylene tubes), 124**
- petit mal seizures, 238**
- pH regulation, 85**
- pharmacodynamics, 14**
- pharmacokinetics, 14**
- pharmacology, 13**
  - administering medications, 17
  - adverse effects, 14
  - angiotensin receptor blockers, 29-30
  - angiotensin-converting agents, 18-19
  - antacids, 15
  - anti-infectives, 15, 20-22
  - antianemics, 15
  - anticholinergics, 16
  - anticoagulants, 15, 33-34, 226
  - anticonvulsants, 16
  - antidiarrheals, 15
  - antihistamines, 15
  - antihypertensives, 15, 18-19, 219
  - antipyretics, 15
  - atypical antipsychotics, 267
  - benzodiazepines, 22-23
  - beta adrenergic blockers, 19-20
  - beta blockers, 219
  - Brethine (terbutaline sulfate), 295
  - bronchodilators, 15
  - burns, 108
  - calcium channel blockers, 219
  - cancer, 144-145
  - cardiovascular disorders, 229
  - chemical names, 18
  - cholesterol-lowering agents, 28-29

- cox 2 enzyme blockers, 30-31
- diuretics, 15, 219
- drug (Hydrocodone) identification, 35
- drug identification, 18, 34
- drug schedules, 36
- ear disorders, 127
- endocrine system disorders, 211
- enteric-coated tablets, 16
- exam prep questions, 37-40
- eye disorders, 123
- gastrointestinal disorders, 171
- generic names, 18
- glucocorticoids, 25
- hematopoietic disorders, 77
- herbals, 35-36
- histamine 2 antagonists, 31-32
- laxatives, 15
- miotics, 16
- musculoskeletal disorders, 193
- mydriatics, 16
- narcotics/analgesics, 16
- neurological disorders, 251
- nitroglycerine, 223
- obstetric clients, 306
- pediatric clients, 343
- pharmacodynamics, 14
- pharmacokinetics, 14
- pharmacologic management of labor, 301-302
- pharmacotherapeutics, 14
- phenothiazines, 24
- pregnancy categories, 36
- protease inhibitors, 26-27
- proton pump inhibitors, 32-33
- psychiatric disorders, 277
- renal/genitourinary disorders, 65
- respiratory disorders, 50-51
- spansules, 16
- time-released drugs, 16
- trade names, 18
- trough drug levels, 22
- Yutopar (ritodrine), 295
- pharmacotherapeutics, 14**
- phases of labor, 296**
- Phenergan, 14**
- phenothiazines, 24**
- phenylketonuria (PKU), 327-328**
- phlebostatic axis, 106**
- phobic disorders, 261**
- phosphorus, normal electrolyte values, 92**
- photorefractive keratotomy (PRK), 121**
- physical therapy for total knee replacement, 189**
- physiologic jaundice, 304**
- pinching, 356**
- pink puffers, 44**
- Pitocin, 298**
- pituitary disorders, 200-201**
- PKU (phenylketonuria), 327-328**
- Placenta Previa, 291**
- placentas, 291**
- pleurisy, 47-48**
- plumbism, 341**
- pneumonia, 46-47**
- POAG (primary open-angle glaucoma), 117**
- poisoning**
  - acetaminophen overdose, 341
  - iron, 342
  - lead, 341
  - salicylate overdose, 341
- polycythemia vera, 76**
- polydipsia, 208**
- polyethylene tubes (PE), 124**
- polyphagia, 208**
- polyuria, 208**
- position, 296**
- positional congenital clubfoot, 323**
- positive schizophrenia, 266**
- positive signs of pregnancy, 285**

**post-traumatic stress disorder (PTSD), 259**

**postpartum care, 302**

**potassium, normal electrolyte values, 91**

**precipitate delivery, 297**

**preeclampsia, 289-290**

**pregnancy**

abortions, 288

abruptio placenta, 291

complications, 287

contraception, 304-305

cord prolapse, 291

DIC (disseminated intravascular coagulation), 290

labor

definition of, 295

dilation, 297

dystocia, 297

effacement, 297

factors influencing, 295-296

fetal lie, 297

fetal monitoring, 298-300

pharmacologic management, 301-302

phases, 296

position, 296

precipitate delivery, 297

prelabor testing, 298

presentation, 296

preterm labor, 295

stages, 296

station, 297

maternal diabetes, 289

maternal infections, 291-294

physiologic jaundice, 304

Placenta Previa, 291

postpartum care, 302

preeclampsia, 289-290

pregnancy categories for drugs, 36

prenatal care

alpha-fetoprotein screening, 285-286

amniocentesis, 286

diagnostic tests, 286

diet and weight maintenance, 285

fetal heart tones, measuring, 287

ultrasonography, 287

Rh incompatibility, 303

signs of

positive signs, 285

presumptive signs, 284

probable signs, 284-285

**pregnancy categories for drugs, 36**

**prelabor testing, 298**

**prematurity, 295**

**prenatal care**

alpha-fetoprotein screening, 285-286

amniocentesis, 286

diagnostic tests, 286

diet and weight maintenance, 285

fetal heart tones, measuring, 287

ultrasonography, 287

**preparing for NCLEX-PN exam**

CAT (Computer Adaptive Test), 4-5

test-taking strategies, 5-8

**presbycusis, 125**

**presbyopia, 121**

**preschoolers, growth and development, 315-316**

**presentation, 296**

**presumptive signs of pregnancy, 284**

**preterm labor, 295**

**prevention of cancer, 137**

**primary hypertension, 218**

**primary open-angle glaucoma (POAG), 117**

**PRK (photorefractive keratotomy), 121**

**probable signs of pregnancy, 284-285**

**prodromal stage (hepatitis), 162-163**

**proliferative diabetic retinopathy, 119**

**protease inhibitors, 26-27**

**Protestant clients, 360**

proton pump inhibitors

**proton pump inhibitors, 32-33**

**psychiatric disorders, 257-258. *See also* neurological disorders**

anxiety-related disorders, 258

DID (dissociative identity disorder), 259-260

GAD (generalized anxiety disorder), 258-259

OCD (obsessive-compulsive disorder), 261-262

panic disorder, 260-261

phobic disorders, 261

PTSD (post-traumatic stress disorder), 259

somatoform disorder, 260

diagnostic tests, 277

disorders of childhood and adolescence

ADHD (attention deficit hyperactive disorder), 276

conduct disorder, 275

eating disorders, 276

oppositional defiant disorder, 275

DSM-IV-TR (Diagnostic and Statistical Manual of Mental Disorders), 258

exam prep questions, 278-281

personality disorders

antisocial personality disorder, 264

avoidant personality disorder, 265

borderline personality disorder, 264

client management, 265

dependent personality disorder, 265

histrionic personality disorder, 263

narcissistic personality disorder, 263

obsessive-compulsive personality disorder, 265

paranoid personality disorder, 262

schizoid personality disorder, 263

schizotypal personality disorder, 263

pharmacology categories, 277

psychotic disorders, 266

bipolar disorders, 268

major depression, 269-270

schizophrenia, 266-268

substance abuse

alcoholism, 270-271

cannabis, 274-275

hallucinogens, 274

opiates, 273

sedative-hypnotics, 273

stimulants, 274

**psychological care, burn patients, 102**

**psychotic disorders**

bipolar disorders

acute mania, 268

major depression, 269-270

schizophrenia, 266-268

**PTSD (post-traumatic stress disorder), 259**

**PTT (partial thromoplastin time), 226**

**pudendal blocks, 301**

**pulmonary disorders. *See* respiratory disorders**

**pulmonary embolus, 42-44**

**purine, 185**

**pyloric stenosis, 334**

## Q - R

**RA (rheumatoid arthritis), 186-187**

**radial keratotomy (RK), 121**

**radiation therapy, 137-138**

**Raynaud's Syndrome, 227**

**reading questions carefully, 6**

**Recombivax, 160**

**red urine, 64**

**refractory errors, 121-122**

**regional enteritis (Crohn's disease), 155**

**regulation of pH, 85**

**regulatory laws, 369**



**rehabilitative phase (burns), 108**

**religious beliefs, 360-361.**

*See also* cultural practices

**renal transplants, 61**

**renal/genitourinary disorders, 57**

acute glomerulonephritis, 58-59

bladder cancer, 64-65

BPH (benign prostatic hyperplasia),  
63

chronic glomerulonephritis, 59

diagnostic tests, 65

ESRD (end stage renal disease),  
60-61

exam prep questions, 66-69

nephrotic syndrome, 61-62

pharmacology categories, 65

urinary calculi, 62

UTIs (urinary tract infections), 62-63

**Respigam, 333**

**respiratory acidosis, 87-88**

**respiratory alkalosis, 90-91**

**respiratory disorders, 41, 328**

acute epiglottitis, 331-332

acute otitis media (AOM), 329

acute respiratory failure

ARDS (acute respiratory distress  
syndrome), 42

pulmonary embolus, 42-44

acute respiratory infections

pleurisy, 47-48

pneumonia, 46-47

TB (tuberculosis), 48

bronchiolitis, 332-333

COPD (chronic obstructive  
pulmonary disease)

asthma, 45

chronic bronchitis, 44

emphysema, 44-45

cystic fibrosis, 333

diagnostic tests, 50

diagnostic tests, 50

emerging infections, 48-49

exam prep questions, 52-55

LTB (laryngotracheobronchitis), 331

pharmacology, 50-51

tonsillitis, 330

**restraints, 374**

**retinal disorders**

diabetic retinopathy, 119-120

hypertensive retinopathy, 119

macular degeneration, 120

retinal detachment, 120

**Rh incompatibility, 303**

**rhabdomyolysis, 28**

**rheumatic fever, 335-336**

**rheumatoid arthritis, 186-187**

**ribavirin, 332**

**right occiput anterior (ROA), 296**

**rights of administering medications, 17**

**rights of clients, 370**

**risk factors for cancer, 135-136**

**risperidone, 267**

**ritodrine, 295**

**RK (radial keratotomy), 121**

**ROA (right occiput anterior), 296**

**RSV-IGIV (Respigam), 333**

**Rule of Nines, 100**

**Russian Orthodox clients, 361**

## S

**SA (sinoatrial) node, 219**

**saccular aneurysm, 227**

**Safe Effective Care, 373-376**

**salicylate overdose, 341**

**salmonella, 170**

**santero/santera, 352**

**sarcoma, 134, 340**

**SARS (Severe Acute Respiratory  
Syndrome), 49**

**Schedule I (drugs), 36**

**Schedule II (drugs), 36**

**Schedule III (drugs), 36**

**Schedule IV (drugs), 36**

**Schedule V (drugs), 36**

**Schilling test, 76**

**schizoid personality disorder, 263**

**schizophrenia, 266-268**

**schizotypal personality disorder, 263**

**school age children, growth and development, 316**

**SCI (spinal cord injuries), 246-248**

**scoliosis, 338-339**

**second degree burns, 99-100**

**second-degree heart block, 220**

**secondary glaucoma, 118**

**secondary hypertension, 218**

**sedative-hypnotic abuse, 273**

**sedatives**

benzodiazepines, 22-23

pregnancy, 301

**seizures**

causes of, 236

generalized seizures, 236

absence, 238

tonic-clonic, 236-237

partial seizures, 238

treatment, 238

**selective serotonin reuptake inhibitors (SSRI), 270**

**self-exams (cancer), 137**

**sensorineural disorders, 115**

diagnostic tests, 126

ear trauma, 126

exam prep questions, 128-130

hearing loss, 126

intraocular disorders

cataracts, 116-117

glaucoma, 117-119

Meniere's disease, 124-125

otitis externa, 124

otitis media, 124

otosclerosis, 125

pharmacology categories, 123, 127

presbycusis, 125

refractory errors, 121-122

retinal disorders

diabetic retinopathy, 119-120

hypertensive retinopathy, 119

macular degeneration, 120

retinal detachment, 120

traumatic injuries, 122

visual tests, 123

**sensorineural hearing loss, 125**

**septic abortions, 288**

**septic emboli, 44**

**set pacemakers, 221**

**seven rights of administering medications, 17**

**Severe Acute Respiratory Syndrome (SARS), 49**

**shamans, 354**

**sickle cell anemia, 74**

**side effects of medications**

angiotensin-converting agents, 19

anti-infectives, 21

benzodiazepines, 23

beta adrenergic blockers, 20

cholesterol-lowering agents, 28

glucocorticoids, 25-26

phenothiazines, 24

protease inhibitors, 27

proton pump inhibitors, 33

**Sikh clients, 356, 361**

**simple fractures, 178**

**simple partial seizures, 238**

**sinoatrial (SA) node, 219**

**skeletal traction, 179**

**skin traction, 179**

**sodium, normal electrolyte values, 91**

**solymigratory arthritis, 336**

**somatoform disorder, 260**

- Somogyi effect, 210**  
**southern belle syndrome, 263**  
**spansules, 16**  
**specific determiners, 7**  
**spina bifida, 323**  
**spinal (subarachnoid) anesthesia, 301**  
**spinal accessory nerve assessment, 244**  
**spinal cord injuries, 246-248**  
**spinal/epidural narcotics, 302**  
**splitting, 264**  
**SSRI (selective serotonin reuptake inhibitors), 270**  
**stigmatism, 121**  
**St. John's wort, 36**  
**Stage 1 withdrawal, 271**  
**Stage 2 withdrawal, 271**  
**Stage 3 withdrawal, 271**  
**Stage 4 withdrawal, 271**  
**stages of labor, 296**  
**stairs crutch-walking gait, 191**  
**standard wound dressings, 107**  
**stapes, 125**  
**staphylococcal, 170**  
**statin drugs, 28**  
**station, 297**  
**status epilepticus, 239**  
**statutory laws, 369**  
**sterilization (contraception), 305**  
**stimulant abuse, 274**  
**strategies for successful test-taking, 5-8**  
**streptokinase, 44**  
**string signs, 155**  
**subacute subdural hematomas, 240**  
**subarachnoid (spinal) anesthesia, 301**  
**subcutaneous nodules, 336**  
**subdural hematomas, 239-240**  
**substance abuse**  
     alcoholism, 270-271  
     cannabis, 274-275  
     hallucinogens, 274  
     opiate abuse, 273  
     sedative-hypnotics, 273  
     stimulants, 274  
**suicide prevention, 269**  
**superficial partial thickness (first degree) burns, 99**  
**surgery, cancer, 137**  
**surgical procedures, musculoskeletal**  
     amputations, 189-190  
     fractured hip and hip replacement, 187-188  
     total knee replacement, 188-189  
**susto (fright sickness), 353**  
**Swan-Ganz catheters, 226**  
**swimmer's ear, 124**  
**swing-through crutch-walking gait, 191**  
**Sydeham's chorea, 336**  
**syngeneic transplants, 141**  
**Synthroid, 202**  
**syphilis, 292**  
**systolic pressure, 218**
- 
- T**
- talipes equinovarus (congenital clubfoot), 323**  
**TB (tuberculosis), 48**  
**TBSA (total body surface area), 98-101**  
**TEF (tracheoesophageal fistula), 319**  
**teratologic congenital clubfoot, 323**  
**terbutaline sulfate, 295**  
**test items, 6**  
**test-taking strategies, 5-8**  
**tet attacks, 326**  
**tetracycline, 35, 47**  
**tetralogy of Fallot (TOF), 326-327**  
**thalassemia major, 75**  
**thickness of the burn injuries, 98-100**  
**third-degree heart block, 220**

**Thomas ring splints, 180**  
**threatened abortions, 288**  
**three-point crutch-walking gait, 191**  
**thromboangilitis obliterans, 226**  
**thrombophlebitis, 227**  
**thunderbirds (Native American amulets), 354**  
**thyroid disorders, 201-203**  
**thyroid storms, 203**  
**thyrotrophic hormone, 200**  
**time considerations (cultural)**  
     Arab-Americans, 358  
     Asian-Americans, 356  
     Hispanics/Latinos, 353  
     Native Americans, 354  
**time-released drugs, 16**  
**toddlers, growth and development, 314**  
**TOF (tetralogy of Fallot), 326-327**  
**tong traction, 180, 247**  
**tonic-clonic seizures, 236-237**  
**tonsillitis, 330**  
**torts, 372**  
**total body surface area (TBSA), 98-101**  
**total knee replacement, 188-189**  
**TPN (total parenteral nutrition), 139-140**  
**tracheoesophageal fistula (TEF), 319**  
**traction, 179-180**  
**trade names, 18**  
**traditional healers**  
     Hispanic/Latino, 352  
     Native Americans, 354  
**transplants**  
     bone marrow transplantation, 140-142  
     renal transplants, 61  
**transspenoidal surgery, 201**  
**transurethral prostatectomy (TURP), 63**  
**traumatic injuries**  
     ear trauma, 126  
     eye injuries, 122

**trigeminal nerve assessment, 243**  
**trochlear nerve assessment, 243**  
**trough drug levels, 22, 47**  
**Trousseau's sign, 90, 204**  
**true congenital clubfoot, 323**  
**tubal ligation, 305**  
**tuberculosis (TB), 48**  
**TURP (transurethral prostatectomy), 63**  
**two-point crutch-walking gait, 190**  
**tyrosine, 327**

## U

---

**ulcerative colitis, 156**  
**ulcers**  
     diagnostic tools, 153  
     dumping syndrome, 154  
     duodenal ulcers, 152  
     gastric ulcers, 153  
     treatment, 153-154  
**ultrasonography, 287**  
**umbilical cord prolapse, 291**  
**uncompensated acidosis, 85**  
**uncompensated alkalosis, 85**  
**urinary disorders, 57-58**  
     acute glomerulonephritis, 58-59  
     bladder cancer, 64-65  
     BPH (benign prostatic hyperplasia), 63  
     chronic glomerulonephritis, 59  
     diagnostic tests, 65  
     ESRD (end stage renal disease), 60  
         hemodialysis, 60  
         peritoneal dialysis, 60  
         renal transplants, 61  
     exam prep questions, 66-69  
     nephrotic syndrome, 61-62  
     pharmacology categories, 65  
     urinary calculi, 62  
     UTI (urinary tract infections), 62-63  
**urinary diversions, 64**

---

**V**

---

- V-fib (ventricular fibrillation), 224-226**
- vaccines, H.influenza B conjugate, 331**
- vaginal bleeding, 287**
- vagus nerve assessment, 244**
- variable deceleration of fetal heart tones, 299-300**
- vasectomy, 305**
- ventricular fibrillation (V-fib), 224-226**
- ventricular tachycardia, 224**
- verbal/nonverbal communication**
  - Arab-Americans, 358
  - Asian-Americans, 357
  - Hispanics/Latinos, 353
  - Native Americans, 355
- vestibulocochlear nerve assessment, 243**
- viral pneumonia, 46**
- vitamins, 73**

---

**W**

---

- walkers, 191-192**
- warning signs of cancer, 134**
- Waterhouse-Friderichsen Syndrome, 206**
- wet (exudative) macular degeneration, 120**
- Wilms tumor, 340**
- withdrawal (substance abuse), 271**
- witnessing consent for care, 373**

---

**X - Y - Z**

---

- xenografts, 107**

- yin/yang, 355**
- Yutopar (ritodrine), 295**

- Zyloprim, 185**